CUMULATIVE AUTHOR INDEX
Journal of Economic Entomology
Volume 106, 2013

DOI: http://dx.doi.org/10.1603/JEE/106_2013/author-index

A

Abbasi, S., 115
Abell, K. J., 1145
Abraham, C. M., 1590
Adam, B., 491
Adamczyk, Jr., J. J., 189
Adams, B., 2378
Adaobi, U. P., 2118
Aguirre, L. M., 1871
Ahmad, M., 954
Akhtar, S., 954
Albajes, R., 1659
Albernaz, K. C., 1819
All, J. N., 830, 1669
Allen, K. C., 382
Alm, S. R., 2322
Alonso, E. C., 742
Altosaar, I., 2585
Alvarez, A., 1234
Alvarez, J. M., 597
Alves, A. P., 2195
Alves, L. C., 747
Amarasekare, K. G., 776, 1126
Ameline, A., 473, 683
Amiens-Desneux, E., 1638
Ammar, E.-D., 25
Amundsen, K., 2568
An, X., 142
Anderson, D. J., 2077
Andrés, V. S., 2541
Ang, L.-H., 2167
Anguiano, O. L., 939
Annan, I. B., 597
Appel, A. G., 322
Argilés, R., 1226
Arimoto, M., 661
Arizza, V., 1582
Armstrong, J. S., 189, 1310, 2077
Arthur, F. H., 1064, 1503, 1911
Arzuffi, R., 1373
Asaro, C., 2391
Asiimwe, P., 1260
Athanassiou, C. G., 1064, 1902
Averill, A. L., 1339
Azevedo, L. H., 2183

B

Bacigalupe, L. D., 400
Bagnères, A.-G., 2216
Bagnoli, B., 2349
Bahder, B. W., 482
Bailey, W., 2020
Baird, L., 2568
Bajracharya, N. S., 2208
Baldin, E.L.L., 1036
Baldwin, J. A., 338
Ban, T., 1767
Bankhead-Dronnet, S., 473
Bansal, R., 1479
Barclay, H. J., 1939
Barajas-Ortiz, M., 1559
Barman, A. K., 1209, 2225
Barnes, B. F., 206
Barr, N., 437
Barragán, S., 1553
Barry, J., 2506
Basit, M., 1414
Basualdo, M., 1553
Bauce, É., 1331
Bauer, L. S., 1145
Baxendale, F. P., 2384, 2568
Bechinski, E. J., 2613
Bedascarrasbure, E., 1553
Beers, E. H., 1996
Behle, R. W., 57
Bélanger, S., 1331
Bellamy, D. E., 1155
Beltrà, A., 1486
Bemquerer, M. P., 86
Benassai, D., 633
Benedettelli, S., 229
Benjamin, T., 2027
Bennett, M. M., 1089
Benson, E. P., 2448
Bentley, M., 97
Bernklau, E. J., 2354
Berthiaume, R., 1331
Beuzelin, J. M., 181
Bezerra-Silva, G.C.D., 1772
Biasi, W. B., 2341
Bigio, G., 2280
Bikoba, V., 2341
Biondi, A., 1638, 2310
Bird, J. P., 1747
Bjostad, L. B., 2354
Black, IV, W. C., 959
Blackburn, C. M., 525
Blackford, D. C., 221
Blankenship, E. E., 1274, 2384
Bobadilla, C., 959
Boerma, H. R., 830
Bohnenblust, E., 1250
Bonafos, R., 1699
Bong, L.-J., 2530
Boratynski, T., 437
Böttger, J.A.A., 426
Bourrié, I., 2585
Bracalini, M., 229
Braman, S. K., 1590, 1693
Bray, A. M., 1190
Brayner, F. A., 747
Breining, J., 1250
Breitler, J.-C., 2585
Brennan, S. A., 912
Bressan, A., 16, 1878
Brewbaker, J. L., 1878
Brewer, M. J., 189, 1310, 2077
Brink, J., 1025
Briones, L. M., 400
Brown, K. S., 329, 1395
Brown, M. J., 1029
Brown, Z. S., 366
Bruner, R. F., 2577
Buawangpong, N., 1535
Buchman, J. L., 1964
Buckman, K. A., 499
Buczkowski, G., 347
Bundy, C. S., 426
Buntin, G. D., 1669
Buonaccorsi, J. P., 1339
Burgio, G., 2296
Burkness, E. C., 2151
Bush, S. J., 1979
Buss, E. A., 1747
Buss, L., 2275
Butron, A., 1234
Butrón, A., 2241

C

Caballero, P., 50
Cabanas, D., 16
Caddoux, L., 1838
Cai, L., 142
Cai, Y., 36
Cambridge, J., 1495
Cambron, S. E., 1164
Cameron, R., 597
Campbell, J. F., 499
Campbell, L. R., 2234
Campos, C., 1045
Cao, A., 2241
Cao, G., 2514
Cao, L.-F., 1958
Cao, Y., 419
Caprio, M. A., 382
Caradonna, S., 633
Cardona, C., 1871
Carloni, E., 1574
Carpane, P., 1574
Carrillo, D., 2286
Cartea, M. E., 1134

CUMULATIVE AUTHOR INDEX
Journal of Economic Entomology
Volume 106, 2013

DOI: http://dx.doi.org/10.1603/JEE/106_2013/author-index

A

Abbasi, S., 115
Abell, K. J., 1145
Abraham, C. M., 1590
Adam, B., 491
Adamczyk, Jr., J. J., 189
Adams, B., 2378
Adaobi, U. P., 2118
Aguirre, L. M., 1871
Ahmad, M., 954
Akhtar, S., 954
Albajes, R., 1659
Albernaz, K. C., 1819
All, J. N., 830, 1669
Allen, K. C., 382
Alm, S. R., 2322
Alonso, E. C., 742
Altosaar, I., 2585
Alvarez, A., 1234
Alvarez, J. M., 597
Alves, A. P., 2195
Alves, L. C., 747
Amarasekare, K. G., 776, 1126
Ameline, A., 473, 683
Amiens-Desneux, E., 1638
Ammar, E.-D., 25
Amundsen, K., 2568
An, X., 142
Anderson, D. J., 2077
Andrés, V. S., 2541
Ang, L.-H., 2167
Anguiano, O. L., 939
Annan, I. B., 597
Appel, A. G., 322
Argilés, R., 1226
Arimoto, M., 661
Arizza, V., 1582
Armstrong, J. S., 189, 1310, 2077
Arthur, F. H., 1064, 1503, 1911
Arzuffi, R., 1373
Asaro, C., 2391
Asiimwe, P., 1260
Athanassiou, C. G., 1064, 1902
Averill, A. L., 1339
Azevedo, L. H., 2183

B

Bacigalupe, L. D., 400
Bagnères, A.-G., 2216
Bagnoli, B., 2349
Bahder, B. W., 482
Bailey, W., 2020
Baird, L., 2568
Bajracharya, N. S., 2208
Baldin, E.L.L., 1036
Baldwin, J. A., 338
Ban, T., 1767
Bankhead-Dronnet, S., 473
Bansal, R., 1479
Barclay, H. J., 1939
Barajas-Ortiz, M., 1559
Barman, A. K., 1209, 2225
Barnes, B. F., 206
Barr, N., 437
Barragán, S., 1553
Barry, J., 2506
Basit, M., 1414
Basualdo, M., 1553
Bauce, É., 1331
Bauer, L. S., 1145
Baxendale, F. P., 2384, 2568
Bechinski, E. J., 2613
Bedascarrasbure, E., 1553
Beers, E. H., 1996
Behle, R. W., 57
Bélanger, S., 1331
Bellamy, D. E., 1155
Beltrà, A., 1486
Bemquerer, M. P., 86
Benassai, D., 633
Benedettelli, S., 229
Benjamin, T., 2027
Bennett, M. M., 1089
Benson, E. P., 2448
Bentley, M., 97
Bernklau, E. J., 2354
Berthiaume, R., 1331
Beuzelin, J. M., 181
Bezerra-Silva, G.C.D., 1772
Biasi, W. B., 2341
Bigio, G., 2280
Bikoba, V., 2341
Biondi, A., 1638, 2310
Bird, J. P., 1747
Bjostad, L. B., 2354
Black, IV, W. C., 959
Blackburn, C. M., 525
Blackford, D. C., 221
Blankenship, E. E., 1274, 2384
Bobadilla, C., 959
Boerma, H. R., 830
Bohnenblust, E., 1250
Bonafos, R., 1699
Bong, L.-J., 2530
Boratynski, T., 437
Böttger, J.A.A., 426
Bourrié, I., 2585
Bracalini, M., 229
Braman, S. K., 1590, 1693
Bray, A. M., 1190
Brayner, F. A., 747
Breining, J., 1250
Breitler, J.-C., 2585
Brennan, S. A., 912
Bressan, A., 16, 1878
Brewbaker, J. L., 1878
Brewer, M. J., 189, 1310, 2077
Brink, J., 1025
Briones, L. M., 400
Brown, K. S., 329, 1395
Brown, M. J., 1029
Brown, Z. S., 366
Bruner, R. F., 2577
Buawangpong, N., 1535
Buchman, J. L., 1964
Buckman, K. A., 499
Buczkowski, G., 347
Bundy, C. S., 426
Buntin, G. D., 1669
Buonaccorsi, J. P., 1339
Burgio, G., 2296
Burkness, E. C., 2151
Bush, S. J., 1979
Buss, E. A., 1747
Buss, L., 2275
Butron, A., 1234
Butrón, A., 2241

C

Caballero, P., 50
Cabanas, D., 16
Caddoux, L., 1838
Cai, L., 142
Cai, Y., 36
Cambridge, J., 1495
Cambron, S. E., 1164
Cameron, R., 597
Campbell, J. F., 499
Campbell, L. R., 2234
Campos, C., 1045
Cao, A., 2241
Cao, G., 2514
Cao, L.-F., 1958
Cao, Y., 419
Caprio, M. A., 382
Caradonna, S., 633
Cardona, C., 1871
Carloni, E., 1574
Carpane, P., 1574
Carrillo, D., 2286
Cartea, M. E., 1134

Carvalho, L. A., 727
Casanoves, F., 2027
Casavecchia, S., 2296
Castañera, P., 2541
Castillo, V. P., 1794
Castle, S. J., 1404
Catchot, A. L., 807, 862, 1430, 2378
Cavalcanti, M.G.S., 747
Chailleux, A., 2310
Chandran, P., 2234
Chang, X., 2548
Chantawannakul, P., 1535
Chaudhury, M. F., 1927
Che, W., 1855
Chen, C., 107
Chen, C.-N., 837
Chen, G., 924
Chen, J., 1832
Chen, L., 142, 1473
Chen, M., 1103
Chen, X., 136, 2630
Chen, X.-M., 1628
Chen, Y., 10, 800, 1053, 1653
Chen, Z.-Z., 1
Cheng, C.-H., 837
Cheng, J., 513
Cheng, W.-N., 2043
Cheng, Z., 36
Cheng, Z.-Q., 1011
Chevrier, C., 641
Chi, H., 1
Choate, B., 716
Christian, K., 299
Cichón, L. B., 939
Clark, C. A., 1566
Clarke, A. R., 695
Claudino, D., 86
Clavet, C., 2322
Clifton, E. H., 622
Cline, A. R., 305
Clodius, M., 786
Cloyd, R. A., 247
Cluck, D. R., 221
Coates, B. S., 2467
Coleman, T. W., 235
Comas, J., 1659
Cook, D., 862, 1430, 2378
Cook, L. G., 2391
Coombs, J. J., 1025
Cooper, M. L., 1716
Cooper, R., 347, 1802
Cooper, W. R., 124
Coots, C., 2399
Correia, A. A., 747
Costello, S., 221
Cottrell, T. E., 257
Coulette, Q., 683
Couty, A., 683
Coyle, D. R., 2391
Cram, M. M., 2391
Crane, J. H., 2286
Creffield, J. W., 329, 1395
Croci, F., 229
Crook, D. J., 2407
Crowe, C. M., 206
Cui, J.-X., 2043
Culbreath, A. K., 587
Curole, J. P., 329

D

Daane, K. M., 482, 1716
Dai, H.-G., 1240
Daigle, L.-F., 1331
Damos, P., 675
Daugovish, O., 80
Davis, A. R., 2055
Davis, J. A., 1566
DeFoliart, L. S., 669
de Guzman, L. I., 566
de la Peña, 408
Delétré, E., 1699
de Mendoza, A. H., 891
Denholm, I., 1414
de Santi, M. P., 1395
Desneux, N., 1638, 2310
Deutschlander, M. E., 2602
Diaz, B. M., 891
Dickey, A. M., 1355
Dickinson, K. L., 366
Dively, G. P., 1317, 2151
Doddala, P.R.C., 899
Dodds, D., 2378
Dodds, K. J., 1684
Dong, B., 1887
Donze, T., 2568
Dorado, J., 2327
Dou, W., 970
Douches, D. S., 1025
Doughty, H. B., 283
Douglas, J., 1517
Driesche, R. V., 1145
Drummond, F. A., 716
Duan, J. J., 1145, 2101
Duan, Y., 2043
Duerr, D. A., 206
Dunbar, M. W., 168, 622
Dunning, L. B., 221
Dupont, S., 2216
Dyer, J. M., 2467

E

Eager, T., 221
Eaton, M., 2433
Ebert, P. R., 2259
Edwards, K. T., 382
Egan, S. P., 1511
Eger, J. E., 912
Eglitis, A., 1684
Eigenbrode, S. D., 2613
Ellsworth, P. C., 1260
Elvira, S., 50
El-Wakkad, M. F., 1608
Enkegaard, A., 2371
Estes, R. E., 1941, 2195
Estévez, A., 1602
Evangelista, D., 2275
Evans, T. A., 329, 1395
Evenden, M. L., 844
Evergetis, E., 1902

F

Fadhl, S., 641
Fail, J., 2602
Falibene, A., 929
Fan, J., 36
Farsani, N. S., 115
Fathi, S.A.A., 1018
Favela, S., 1004
Feder, J. L., 1511
Felcher, K. J., 1025
Feldlaufer, M. F., 988
Fereres, A., 891
Fernández-Quintanilla, C., 2327
Fettig, C. J., 1684
Fielding, D. J., 669
Figueroa, C. C., 400
Fiuza, L. M., 2585
Fleischer, S., 1250
Flint, M. L., 235
Flores, A. E., 959, 1004
Flor-Weiler, L. B., 57
Follett, P. A., 653, 2020, 2035
Fontaine, S., 1838
Fontenot, E. A., 1503, 2336
Fontes, E.M.G., 86
Forim, M. R., 1772
Foroughbakhch, R., 1004
Foster, N., 437
Foster, S. P., 400, 855
Frake, A. M., 566
Francese, J. A., 2407, 2415
Frank, D. L., 2506
Franz, G., 995
Fraser, I., 2407
Freeman, J. H., 283
Freitag, J. A., 1920
Frewin, A. J., 552
Freytag, E. D., 329, 1395
Frost, E. H., 558
Fu, Q., 1887
Fu, R. S., 43
Fuentes-Contreras, E., 400
Fujisaki, I., 311
Fukumoto, C., 463
Funayama, K., 2141
Furlan, P., 633

G

Galmarini, C. R., 1386
Ganbalani, G. N., 1018
Gandhi, K.J.K., 206, 2391
Gantet, P., 2585
Gao, C., 1832, 2190
Gao, S., 1294
Garcerá, C., 2126

García, C., 1553
Garcia-Marí, F., 1486
Gardescu, S., 874
Garnas, J. R., 1979
Garrido, S.A.S., 939
Garrido-Jurado, I., 1118, 1602
Garzón-Tiznado, J. A., 1559
Gassmann, A. J., 168, 622, 1941, 2195, 2577
Gaussoin, R. E., 2384
Gautam, B. K., 1812, 2630
Gautam, S. G., 491
Ge, L.-Q., 196
Geisert, R. W., 1619
Gelbič, I., 1075, 1098
Gil, R. A., 1386
Giles, K. L., 491
Gill, B. S., 1911
Godfrey, L., 1707
Goenaga, R., 305
Gondhalekar, A. D., 945
Gong, Z.-J., 2043
Gonçalves, G.G.A., 747
Gore, J., 807, 862, 1430, 2378
Gorria, N., 50
Gotoh, T., 463
Gradish, A. E., 552
Grafton-Cardwell, E. E., 2126, 2341
Grant, J., 2399
Graves, A. D., 235
Gray, M. E., 1941, 2195
Greene, J. K., 1676, 2448
Greenlee, K. J., 1081, 1089
Greenplate, J., 1525
Grenkow, L., 2557
Gross, D., 1440
Groves, R. L., 2491
Gu, H., 883
Guan, X., 1075, 1098
Guerfali, M. M., 641
Guiderdoni, E., 2585
Gullan, P. J., 2391
Guo, J.-Y., 1379
Guo, T.-T., 1
Guo, W., 1112
Guo, X.-J., 2621
Gut, L. J., 2144
Gutierrez, S. M., 959

H

Hagerty, A. M., 669
Hague, J.R.B., 329
Hajek, A. E., 874
Hall, D. G., 25
Hall, P. M., 1355
Hallett, R. H., 267
Hallman, G. J., 525, 1173, 1608, 2336
Hamacek, E. L., 883
Hamby, K. A., 1365, 1932
Hamden, H., 641
Hamilton, G. C., 1495
Hammon, R. W., 1029
Hampton, E., 2322
Han, P., 2310
Han, R., 1112, 1863
Hancock, J., 1440
Hanson, S. F., 426
Hanula, J. L., 1693
Hao, Y.-N., 2043
Harding, C., 786
Hardy, G., 1240
Harney, M., 1979
Haroutounian, S. A., 1902
Harrison, Jr., H. F., 1456
Hashiyama, A., 690
Haverty, M. I., 338
Haviland, D. R., 1986
Hayes, D. B., 1324
He, J., 1887
He, K. L., 2548
He, W.-B., 1446
He, Y., 1832
Head, G. P., 1525, 1819
Hébert, C., 1331
Hein, G. L., 1274
Henderson, G., 1812, 2630
Hendrix, III, W. H., 823
Heng-Moss, T. M., 1036, 2384, 2568
Henne, D. C., 1964
Hepburn, H. R., 1958
Herbert, Jr., D. A., 1317
Hibbard, B. E., 2354, 2506
Hidalgo, E., 2027
Higashi, C.H.V., 16, 1878
Higdon, M. L., 2506
Hillier, N. K., 558
Hinkle, N. C., 1590
Hinomoto, N., 463
Hodgson, E. W., 2577
Hoffmann, A. A., 756
Hoffmann, B. D., 1545
Hofstetter, R. W., 1684
Hollingsworth, R. G., 653
Holmstrom, K., 1495
Horton, D. R., 1996
Howell, A. D., 80
Hsu, Y.-C., 408
Hu, F., 970
Hu, F.-L., 1958
Hu, X.-S., 1894
Hu, Y.-Q., 1958
Hu, Z.-Q., 1894
Huang, E., 1098
Huang, J., 2144
Huang, S.-H., 837
Huang, T., 1075, 1098
Huang, T.-I., 1726
Huang, Y. H., 43
Huang, Z., 1075, 1098
Huang, Z. Y., 1653
Hulasare, R., 1173
Hulcr, J., 2093
Hunt, T. E., 1036, 1274, 2384
Huo, X., 360
Huo, X. B., 43
Hurley, B. P., 1979
Husen, T. J., 1788
Huseth, A. S., 2491
Hutchison, W. D., 2151

I

Ibargutxi, M. A., 50
Ide, Y., 536
Iline, I. I., 2006
Ingerson-Mahar, J., 1495
Inoue, H., 536
Ioriatti, C., 2349
Irish, B., 305
Isaacs, R., 905
Isidoro, N., 604, 2296
Iwanami, T., 536

J

Jaal, Z., 2530
Jacas, J. A., 1226
Jackson, D. M., 1456
Jackson, M. A., 57, 257, 622
James, J., 648
Jang, E. B., 653, 727
Jankielsohn, A., 2595
Jaronski, S. T., 622
Jayas, D. S., 2247
Jean, G. S., 1511
Jenkins, D. A., 305
Jenkins, S., 756
Jessup, A., 695
Jessup, A. J., 1608
Jian, F., 2247
Jiang, J., 142
Jiang, L.-B., 196
Jiang, Y.-L., 2043
Jiménez-Pérez, A., 1373
Jin, T., 353
Jing, X., 43
Johnson, R. M., 1952
Jones, C. L., 2208
Jones, M. I., 235
Jorgensen, C. L., 221
Josens, R., 929
Juan-Blasco, M., 1226, 2541

K

Kadirvel, P., 408
Kafle, L., 131
Kamal, A., 1699
Kamble, S. T., 1788
Kang, D. M., 43
Kang, J. K., 2473
Kard, B., 1395
Kard, B. M., 329
Katoh, H., 536
Katsoula, A.-M., 1902
Kaur, R., 2259

Kavallieratos, N. G., 1064, 1902
Keller, M. A., 1894
Kells, S. A., 1920, 2433
Kemp, W. P., 1081, 1089
Kerns, D., 1430
Keskin, S., 1058
Khakame, S. K., 1423
Khan, R. A., 2177
Khan, S. A., 2234
Kheradmand, K., 115
Killiny, N., 393
Kim, T., 2428
Kinjo, H., 1767
Kitsis, P., 1848
Knutson, A. E., 1045
Kong, L., 1473
Kopittke, R. A., 883
Kramer, M., 988
Kramer, R. A., 366
Krause, C. R., 159
Kuah, M.-K., 2167
Kuchiki, F., 536
Kuhar, T. P., 283, 1317
Kunimi, Y., 1767
Kurihara, J., 690
Kučerová, Z., 419

L

Labédan, M., 2216
Labrie, J., 1331
Laguna, I., 1574
Lambdin, P., 2399
Lance, D. R., 2407
Landi, L., 604
Lang, E. B., 597
Langemeier, M. R., 1503
Langor, D. W., 1684
Lanka, S. K., 181
Larson, K., 2101
Lasue, P., 683
Lavine, L. S., 2424
Lee, C.-Y., 1395, 2167, 2530
Lee, D.-H., 150
Lee, J. C., 2424
Leighton, S., 338
Lelito, J. P., 1145, 2101
Lenz, M., 329, 1395
Leon, R. D., 437
Leonard, B. R., 1430
León-Sicairos, C. R., 1559
Leopold, R. A., 855
Leskey, T. C., 150
Lévy, J., 1440
Lewis, V. R., 338
Li, F., 419, 924, 1473
Li, G., 36
Li, G.-Q., 614
Li, H., 450, 708
Li, H.-F., 311
Li, J. L., 513
Li, J.-M., 1446
Li, K., 1653
Li, L.-L., 1011
Li, S., 2457
Li, T., 2043
Li, W., 1200
Li, X., 2514
Li, Y., 375, 1200
Li, Z., 419
Liang, G., 2514
Liang, Y., 513
Liburd, O. E., 73, 912, 1757
Lichtenberg, E. M., 1535
Lieutier, F., 473
Lim, U. T., 1973
Lin, H., 2457
Lin, Y., 353, 1887
Liu, B., 924
Liu, C., 347, 2522
Liu, D., 1294
Liu, G. G., 2548
Liu, H., 360
Liu, H.-Y., 1825
Liu, J. J., 43
Liu, Q., 10, 360
Liu, Q. Z., 1349
Liu, S.-S., 1446
Liu, S. S., 1613
Liu, T.-X., 1894
Liu, W., 360
Liu, W.-F., 970
Liu, X., 450, 708, 1103
Liu, X.-F., 1894
Liu, Y., 1473
Liu, Y.-B., 1155, 1613, 2267
Liu, Z., 1098
Liu, Z.-D., 1240
Liu, Z. L., 513, 1349
Lloyd, A. C., 883
Long, B., 1331
Lopez, B., 959, 1004
López-Valenzuela, J. A., 1559
Lorenz, G., 1430
Lowery, D. T., 2135
Lozano-Tovar, M. D., 1118
Lu, W., 215
Lu, Y., 353, 360
Lu, Z., 1887
Lü, L., 2522
Luan, X. B., 1349
Lucchi, A., 2349
Luchetti, A., 2216
Lumbierres, B., 1659
Luo, C., 1653
Luo, Y.-Q., 866
Lv, J., 215

M

Ma, C.-S., 2118
Ma, Y., 614
Mainali, B. P., 1973
Maldonado, I. B., 1386
Malvar, R. A., 1234, 2241
Manachini, B., 1582
Maner, M. L., 1693
Mansouri, S. M., 1018
Mantovani, B., 2216
Marcum, D., 1707
Marshall, D., 73
Martín, J. M., 2327
Martin, T., 1699
Martinelli, S., 1819
Martínez-Uña, A., 2327
Mason, K. S., 905
Mastrangelo, T., 1772
Mastro, V. C., 2407, 2415
Matsuda, T., 463
Mayfield, III, A. E., 1684
Mbata, G. N., 2072
McCarville, M. T., 1302
McColl, S., 756
McCullough, D. G., 1324
McFarlane, A. D., 552
McKenzie, C. L., 1355
McNeil, J. N., 1648
McNeill, M. R., 2006
Mech, A. M., 2391
Medina, R. F., 2225
Meeus, I., 277
Meikle, W. G., 648
Meinke, L. J., 1619, 1941, 2195
Mekala, K. D., 1045
Melika, G., 1747
Men, X.-Y., 1011
Menezes, V. G., 2585
Meng, F., 360
Mercado, R., 959
Merlin, B. L., 1819
Merrill, S. C., 2084
Meyer, L. D., 277
Meynard, D., 2585
Meza, J. S., 735
Mian, M.A.R., 1479
Miao, J., 2043
Michel, A. P., 1479
Micoud, A., 1838
Mikac, K. M., 1517
Miles, J. W., 1871
Millar, J. G., 482
Miller, D. R., 206, 1684
Miller, J. R., 2144
Minor, M. A., 899
Minuz, R. L., 2296
Miranda, J. E., 86
Mitcham, E. J., 2341
Moar, W., 1525
Moltó, E., 2126
Mommaerts, V., 277
Montagna, C. M., 939
Moraes, G. J., 2183
Morales, I., 891
Mori, B. A., 844
Morin, V., 2433
Morrison, III, W. R., 1218
Morse, J. G., 64, 1986
Mostafa, A. M., 2135

Moyseenko, J. J., 159, 1190
Mu, L.-L., 614
Mulder, P. G., 1071
Muñoz, D., 50
Munson, A. S., 221, 1684
Munyaneza, J. E., 1964
Musser, F. R., 382, 807, 862, 2378
Mutitu, E. K., 1979
Myers, S. W., 1155, 1608, 1613, 2336

N

Naidu, R. A., 482
Nakai, M., 1767
Nansen, C., 648
Naranjo, S. E., 1260
Naseri, B., 1018
Navarro-Llopis, V., 1739
Nazni, W. A., 2167
Nebreda, M., 891
Nechols, J. R., 1503
Neoh, K.-B., 2530
Neven, L. G., 520
Ngouajio, M., 1699
Ni, H., 450
Nicholls, J. A., 1747
Nicholson, S. J., 1029
Nicolini, G., 2349
Nielsen, A. L., 1495
Nolting, S. P., 823
Nomura, M., 690
Novoselov, M., 2006
Nyoike, T. W., 1757

O

Obenland, D., 1155
Obregón, O., 437
Oesterle, N., 426
Oetting, R. D., 1590
Oliveira, J. V., 747
Oliveira, J.V.D., 2585
Oliver, J. B., 159, 289, 1190, 1780
Olmstead, D. L., 2151
Olson, J. F., 2433
Omoto, C., 1819
O'Neal, M. E., 1302
Onstad, D. W., 1103, 2473
Opit, G. P., 419, 491, 2208
Ordas, B., 1234
Orozco, D., 735
Ortego, F., 1226
Ortiz, A., 1602
Osakabe, M., 661
Osborne, L. S., 1355
Ottea, J. A., 181
Ouyang, Y., 2126, 2341
Owens, D. R., 1317
Ozkaya, H., 1058

P

Pakyari, H., 2371
Palix, R., 1699
Palumbo, J. C., 1726
Pan, H., 924
Pang, Y., 36
Panzavolta, T., 229
Pappu, H. R., 1964
Paradell, S., 1574
Parajulee, M. N., 1209, 2225
Parker, A. G., 525
Parker, R. D., 189, 1310
Patten, K., 2424
Paula, D. P., 86
Paula-Moraes, S., 1274
Payton, M. E., 1071, 1173
Peairs, F. B., 2084
Pecci, M.P.G., 1574
Peck, D. C., 1463
Peck, S. L., 1940
Pedroso, G. M., 2341
Peña, J. E., 2286
Peng, B., 360
Peng, J., 1473
Peng, R., 299
Peng, Y. F., 2548
Peralta, I. E., 1386
Peraza-Garay, F., 1559
Percel, E. G., 1952
Perdereau, E., 2216
Perring, T. M., 1726
Petersen, K., 1089
Petersen, M. J., 1463
Pett, W., 1025
Pettis, J. S., 1535
Petzold-Maxwell, J. L., 622, 1941, 2195
Peverieri, G. S., 633
Pfeiffer, D. G., 283
Phillips, C. B., 2006
Phillips, T. W., 1173
Pierson, E., 1440
Pierson, L. M., 1036
Pieterse, W., 1179
Pinto, L.M.N., 2585
Pires, C.S.S., 86
Pittendrigh, B. R., 2473
Pitts-Singer, T. L., 543, 576
Ploetz, R., 2093
Pointeau, S., 473
Poland, T. M., 1684
Ponce, G., 959, 1004
Pons, X., 1659
Porter, P., 648
Portillo, H. E., 597
Pozo-Valdivia, A.I.D., 533
Prabhaker, N., 1404
Prasifka, P. L., 823
Pratiwi, R., 2364
Primo, J., 1739
Prochaska, T. J., 1036
Progar, R. A., 221
Pupin, F., 2341
Puterka, G. J., 1029

Q

Qi, S., 136
Qian, K., 979
Qin, A.-Z., 215
Qin, J.-L., 866
Qin, M., 419
Qin, X., 10
Qiu, G.-S., 2118
Qiu, S., 1075
Qiu, X., 1863
Quesada-Moraga, E., 1118, 1602
Quintero-Fong, J. L., 735

R

Raffa, K. F., 1684
Rajamohan, A., 855
Rambaud, C., 683
Ramm, C., 2568
Ramsey, A., 727
Ranger, C. M., 159, 289, 1190, 1780
Rashid, M., 2177
Ratnieks, F.L.W., 2280
Ravindran, A., 1440
Razmjou, J., 1018
Reay-Jones, F.P.F., 1676, 2448
Reding, M. E., 159, 289, 1190, 1780
Reed, D. A., 1726
Reese, J. C., 1036, 2234, 2384
Reisig, D. D., 533, 1317
Ren, D., 360
Ren, L.-L., 1240
Requintina, M., 2322
Reuter, C., 437
Revilla, P., 1234
Reyes-Moreno, C., 1559
Reynolds, H. L., 97
Rhea, R., 2399
Rhodes, E. M., 912
Ribeiro, A.C.J., 86
Richards, N. K., 2006
Rietz, M. L., 2407, 2415
Riley, D. G., 587
Rinderer, T. E., 566
Rinehart, J. P., 855, 1081, 1089
Rinella, M. J., 221
Riolo, P., 604, 2296
Rodríguez, E., 1553
Rogers, D. J., 899
Roland, Sr., T. A., 1395
Rose, R., 1535
Rosenheim, J. A., 1286
Roth, G., 1250
Roubos, C. R., 73, 905
Roush, R., 1103
Roversi, P. F., 633
Roy, L., 1838
Ruiz-Arce, R., 437
Ruíz-Torres, M., 1602
Rule, D. M., 823
Rust, M. K., 2428

S

Saathoff, A., 2568
Sabater-Muñoz, B., 1226, 2541
Saccaggi, D. L., 1179
Sadof, C., 2027
Saeed, S., 1414
Sahri, M. H., 1794
Saidi, M., 641
Sajap, A. S., 1794
Saleem, M. A., 1414
Sallé, A., 473
Sampson, B. J., 73
Samuel-Foo, M., 830
Sansone, C. G., 2225
Santiago, R., 2241
Santolamazza-Carbone, S., 1134
Santos, D.Y.A.C., 742
Sappington, T. W., 2467
Saran, R. K., 945
Satoh, M., 661
Schapaugh, W., 2234
Scharf, M. E., 945
Scherer, C. W., 945
Schillaci, D., 1582
Schleip, E. M., 2084
Schlipalius, D. I., 2259
Schmidt, R. A., 1996
Schultz, P. B., 283, 289, 1780
Schürch, R., 2280
Schutze, M. K., 695
Scott, S. J., 2126
Scott, W. S., 862
Scott-Dupree, C. D., 552
Scown, D. K., 329
Sears, M. K., 267
Sehgal, B., 1911
Seiter, N. J., 1676, 2448
Selfa, J., 1134
Sengoda, V. G., 1964
Seybold, S. J., 235, 800
Shah, M., 1414
Shao, S.-X., 1628
Shapiro-Ilan, D. I., 257, 2072
Shatters, Jr., R. G., 1355
Shearer, P. W., 776, 1126
Shelton, A. M., 1103, 1525, 2151, 2602
Shelton, T. G., 2160
Shen, J., 1832
Shen, Z. C., 2548
Shi, A.-J., 1
Shi, P., 107
Shi, T., 1855
Shi, W. P., 1349
Shields, E. J., 648, 2195
Shih, C. J., 131
Shrestha, A., 587
Shu-Chien, A. C., 2167
Shufran, A. A., 1071
Shufran, K. A., 1071
Shupe, T. F., 329
Shutler, D., 558
Siegfried, B. D., 2384
Silva, A. X., 400
Silva, B. G., 959
Silva, M. A., 1772
Silva, M. F., 1386
Simmons, G. S., 1155, 1613
Simon, J.-C., 1838
Simon, S., 1699
Sims, S. R., 979
Singh, N., 347, 1802
Singleton, M. E., 2424
Siri, N., 1648
Siscaro, G., 1638
Situmorang, J., 2364
Skoda, S. R., 1927
Slippers, B., 1979
Smagghe, G., 277
Smirle, M. J., 2135
Smith, A. A., 97
Smith, C. W., 1045
Smith, D., 883
Smith, J. A., 2093
Smith, J. F., 807
Smith, W. R., 329
Snoddy, E. T., 322
Snook, K., 2035
Soengas, P., 1134
Soesilohadi, R.C.H., 2364
Sola, F., 929
Solana, H., 1553
Soleño, J., 939
Solís, E., 735
Somana, W., 1535
Song, M., 136
Sornnuwat, Y., 1395
Soroka, J., 2557
Sotelo, G., 1871
Soto, A., 1486
Soto, V. C., 1386
Sparks, A. N., 2151
Spence, D. J., 2093
Spencer, J. L., 1517
Spurgeon, D. W., 124
Srinivasan, R., 408, 587
Stamm, M. D., 2384
Stanghellini, M. E., 1986
Steckel, S., 816, 2048
Steed, B., 221
Steffan, S. A., 2424
Stejskal, V., 419
Stelinski, L. L., 393, 2093
Stewart, S. D., 816, 1430, 2048
Stokes, B., 648
Stokes, B. S., 2613
Stone, G. N., 1747
Storer, N. P., 823
Stout, M. J., 181
Stringer, S. J., 73
Strong, W. B., 633
Su, F.-C., 408
Su, J., 1832, 2190
Su, N.-Y., 311, 2442
Su, Y.-L., 1446
Subramanyam, B., 499, 1911, 2247
Sujii, E. R., 86
Sukirno, S., 2364
Sukumalanand, P., 1535
Sumarmi, S., 2364
Sumerford, D. V., 1525
Sun, L.-N., 2118
Sun, S., 36
Sun, W. B., 43
Sun, Y., 1653
Sundaraj, S., 587
Swain, A. J., 2259
Szendrei, Z., 1025, 1218

T

Tabone, E., 2310
Tabuchi, R., 338
Tadrous, M. D., 1608
Talley, J., 2208
Tamborindeguy, C., 1440
Tan, X.-L., 2621
Tan, X.-M., 1011
Tang, B., 1098
Tang, Z., 360
Taret, G., 2336
Taylor, D. B., 979
Taylor, P., 2101
Tebbets, S., 1155
Teixeira, Á.A.C., 747
Teixeira, L.A.F., 905
Terreni, P., 229
Tewari, S., 1339
Tholl, D. B., 283
Thompson, S. D., 2195
Throne, J. E., 1064
Tian, M.-Y., 215
Tiberi, R., 229
Timbó, R. V., 86
Tindall, K. V., 2048
Tinsley, N. A., 2195
Tiwari, S., 393
Tollerup, K. E., 1707
Tooker, J., 1250
Toyoshima, G., 690
Trewick, S. A., 899
Tsitsilas, A., 756
Tsunoda, K., 1395
Twigg, P., 2568

U

Uechi, N., 536
Uesugi, R., 661
Ugine, T. A., 874
Ul-Haq, I., 695
Ulrich, K. R., 988
Umina, P. A., 756
Unruh, T. R., 400, 1996
Urbaneja, A., 1226, 2541

V
Vacas, S., 1739
Vacek, D., 437
Valdez-Ortiz, A., 1559
Vanagas, L., 1553
vanEngelsdorp, D., 1535
van Herk, W. G., 786
Varela, L. G., 2349
Vassiliou, V. A., 1848
Velarde-Félix, S., 1559
Velasco, P., 1134
Velonà, A., 2216
Vendramim, J. D., 1772
Vera, M. T., 695
Vernon, R. S., 786
Vilaire, A., 2424
Vongkaluang, C., 1395
Vreysen, M.J.B., 695, 1939, 2336

W
Wall, M., 2020
Wallingford, A. K., 283
Walse, S. S., 1155, 1613
Walsh, D. B., 482, 2424
Walter, A. J., 25
Walter, G. H., 2259
Walter, S. M., 2084
Wan, D.-J., 196
Wan, F.-H., 1379
Wan, P.-J., 614
Wanderley-Teixeira, V., 747
Wang, C., 136, 347, 1802, 1812, 2433, 2522, 2630
Wang, D., 1825, 2234
Wang, H., 1473
Wang, J., 360, 1446
Wang, J.-J., 970
Wang, L., 107
Wang, Q., 142, 215, 1373
Wang, S., 924, 2621
Wang, X., 360, 1423
Wang, X.-C., 1011
Wang, X. J., 43
Wang, X.-W., 1446
Wang, Y., 142
Wang, Y.-M., 1825
Ware, J. L., 2275
Watanabe, S., 16
Watt, T., 2101
Webb, G. A., 1545
Wei, D., 970
Wei, S.-G., 215
Wei, W., 1240
Wei, W.-T., 1958
Welty, C., 2151
Wheeler, T., 648
White, N.D.G., 2247
Wieferich, D. J., 1324
Williams, T., 50
Willmott, A. L., 247
Wilson, R., 1707
Wingfield, M. J., 1979
Wingsanoi, A., 1648
Wist, T. J., 2055
Wood, B. W., 257
Wornoayporn, V., 695
Wosula, E. N., 1566
Wright, R. J., 1274
Wright, S. E., 150
Wu, C., 1075
Wu, D., 360
Wu, G., 1379
Wu, J.-C., 196
Wu, K., 1200, 2514
Wu, M., 2190
Wu, Q., 924
Wu, W.-J., 837
Wu, Y., 375, 1423, 1855
Wu, Y.-J., 866
Wu, Y.-Q., 2043
Wunderlich, L. R., 1716

X
Xie, W., 924
Xin, Z., 1825
Xu, B., 924
Xu, H., 1887
Xu, J., 196, 215
Xu, L., 1075, 1098
Xu, S., 513
Xu, Y.-Y., 1
Xue, F., 107
Xue, M., 1825

Y
Yamamoto, P. T., 2183
Yan, S., 450, 708
Yan, W.-T., 2118
Yan, X., 1112
Yang, Q., 419
Yang, X.-H., 866
Yang, Y., 375, 1855, 1887
Yang, Z.-X., 1628
Yaning, F., 2457
Yao, R., 2190
Ye, G. Y., 2548
Yee, W. L., 1511, 2109
Yeh, B.-H., 1
Yeoh, B.-H., 1395
Yin, W.-D., 2118
Yocum, G. D., 1081, 1089
Yokoyama, V. Y., 1164
Yoshimura, T., 1395
Yousef, M., 1118, 1602
Youssef, N. N., 159, 289, 1190
Yu, D., 1932
Yu, L.-Y., 1
Yu, Y., 1011
Yu, Y.-H., 866
Yuan, W., 1200

Z
Zacharopoulou, A., 995
Zahn, D. K., 64, 1986
Zalom, F. G., 1365, 1932
Zamani, A. A., 115
Zanardi, O. Z., 2183
Zappalà, L., 1638
Zeng, L., 353
Zeng, X., 979
Zepeda, S., 735
Zhang, A., 2522
Zhang, A.-S., 1011
Zhang, F., 43, 2621
Zhang, H., 1349
Zhang, H.-J., 2118
Zhang, J., 10, 450, 708, 1832
Zhang, L., 1075, 1098, 2514
Zhang, Q., 450, 708
Zhang, R., 10
Zhang, S., 2190
Zhang, X. C., 1349
Zhang, Y., 614, 924, 1863, 2177
Zhang, Z., 2190
Zhao, G., 1112
Zhao, H.-Y., 1894
Zhao, N. N., 513, 1349
Zhao, S., 419
Zhao, X., 142
Zhao, Y., 1863
Zheng, F.-Q., 1
Zheng, H.-Q., 1958
Zheng, X., 1887
Zhifeng, X., 2457
Zhong, J.-H., 329, 1395
Zhou, C., 1349
Zhou, H.-X., 1011
Zhou, L., 513
Zhu, J., 708
Zhu, J. J., 979
Zhu, K. Y., 247
Ziegler, D., 2585
Zuehlke, M., 1025
Zukoff, A., 2506
Zungoli, P. A., 2448
Zurowski, C. L., 2135

CUMULATIVE KEY WORD INDEX
Journal of Economic Entomology
Volume 106, 2013

DOI: http://dx.doi.org/10.1603/JEE/106_2013/keyword-index

A

Abagrotis,
  cutworm, 2135
  grape pest management, 2135
  insecticide toxicity, 2135
abamectin,
  acrinathrin, 1848
  bifenazate, 1848
  fenazaquin, 1848
  pirimiphos methyl, 1848
acephate,
  aldicarb, 807
  imidacloprid, 807
  soil-applied insecticides, 807
  thiamethoxam, 807
Acerophagus,
  Ferrisia gilli, 1716
  grape, 1716
  mealybug, 1716
  Vitis, 1716
Acer platanoides,
  distribution, 874
  imidacloprid, 874
  neonicotinoid, 874
  translocation, 874
acoustic emissions,
  drywood termites, 338
  feeding periodicity, 338
  temperature effects, 338
acquisition, 277, 1878
  Bombus terrestris, 277
  life history, 1878
  Metarhizium anisopliae strain F52, 277
  miniature dispenser, 277
  plant virus, 1878
  resistance, 1878
  transmission, 1878
  worker mortality, 277
acquisition access period,
  Liberibacter, 1964
  potato, 1964
  potato psyllid, 1964
  tomato, 1964
acrinathrin,
  abamectin, 1848
  bifenazate, 1848
  fenazaquin, 1848
  pirimiphos methyl, 1848
action threshold,
  degree-day, 1757
  Fragaria ananassa, 1757
  mite-day, 1757
  six-spotted thrip, 1757
actograph,
  Autographa nigrisigna, 690
  flight ability, 690
  flight mill, 690
actuarial mathematics,
  Anarsia lineatella, 675
  demographic parameter, 675
  intrinsic rate of increase, 675
  life table, 675
acute toxicity,
  biological parameter, 1825
  spinosad, 1825
  Spodoptera exigua, 1825
  sublethal effect, 1825
Acyrthosiphon pisum,
  binomial sequential sampling, 2613
  dry pea, 2613
  pea aphid, 2613
  Pisum sativum, 2613
additives,
  Bacillus thuringiensis, 1075
  bioassay, 1075
  biocontrol, 1075
  Plutella xylostella, 1075
adult damage,
  Brachiaria, 1871
  host-plant resistance, 1871
  spittlebug, 1871
adult longevity,
  Cheilomenes sexmaculata development, 2621
  fertility, 2621
  prey consumption, 2621
adult starvation,
  daily reproductive rhythm, 215
  Glenea cantor, 215
  lifetime reproductive pattern, 215
  sexual maturation, 215
Aedes aegypti,
  cross-resistance, 959
  insecticide resistance, 959
  pyrethroid, 959
Aedes albopictus,
  Bacillus thuringiensis, 1098
  ELISA, 1098
  immunohistochemistry, 1098
  receptor binding, 1098
AFLP,
  cotton pest, 2225
  dispersal, 2225
  gene flow, 2225
  population structure, 2225
agricultural landscape,
  crop pollination, 97
  seminatural areas pollinator habitat, 97
  wild bees, 97
Agrilus planipennis, 2101, 2415
  development, 2101
  fluon, 2415
  Fraxinus uhdei, 2101
  green, 2415
  invasive, 2101
  laboratory rearing, 2101
  multifunnel trap, 2415
  purple, 2415
agro-ecosystems,
  invertebrate, 756
  mortality, 756
  nontarget, 756
  pesticide, 756
Alaska,
  Conservation Reserve Program, 669
  grasshopper bait, 669
  high-latitude agriculture, 669
  Insecta, 669
aldicarb,
  acephate, 807
  imidacloprid, 807
  soil-applied insecticides, 807
  thiamethoxam, 807
alfalfa,
  Megachile rotundata, 543
  Megachilidae, 543
  pollination, 543
  reproduction, 543
alfalfa leafcutting bee,
  fluctuating thermal regime, 1081
  insect cold storage, 1081
  Megachile rotundata, 1081
alfalfa-seed production,
  Medicago sativa, 576
  Megachile rotundata, 576
  Megachilidae, 576
  pollination, 576
alfalfa weevil,
  Hypera postica, 426
  phylogenetic, 426

Allium cepa L.,
floral traits, 1386
male sterility, 1386
nectar composition, 1386
seed yield, 1386
Allium sativum,
allyl trisulfide, 1349
Cacopsylla chinensis, 1349
contact toxicity, 1349
diallyl disulfide, 1349
allyl trisulfide,
Allium sativum, 1349
Cacopsylla chinensis, 1349
contact toxicity, 1349
diallyl disulfide, 1349
almond,
boscalid, 1952
diflubenzuron, 1952
organosilicone, 1952
pyraclostrobin, 1952
alphacypermethrin,
Aphidius colemani, 1699
cabbage, 1699
Myzus persicae, 1699
repellent netting, 1699
α-pinene,
ethanol, 1684
ipsdienol, 1684
ipsenol, 1684
kairomone, 1684
alternating generation,
bullet gall, 1747
gall wasp, 1747
Quercus virginiana, 1747
alternative host plant,
biotype, 2595
Diuraphis noxia, 2595
intrinsic rate of increase, 2595
Russian wheat aphid, 2595
ambrosia beetle,
bottle trap, 1780
forecasting activity, 1780
phenological indicator, 1780
trap tree, 1780
amitraz,
honey bee, 558
miticides, 558
tau-fluvalinate, 558
Varroa destructor, 558
Amyelois transitella,
bird damage, 1365
Corvidae, 1365
degree-day, 1365
Prunus dulcis, 1365
survivorship, 1365
Anarsia lineatella,
actuarial mathematics, 675
demographic parameter, 675
intrinsic rate of increase, 675
life table, 675
Anastrepha ludens,
genetic sexing, 735
sexual behavior, 735
sterile insect technique, 735
Tapachula-7, 735
ant, 929, 1545
bait, 929
boric acid, 929
control, 1545
eradication, 1545
feeding, 929
invasive ants, 1545
sodium borate, 929
treatment, 1545
antagonism,
bioassay, 247
horticulture, 247
pest management, 247
synergism, 247
antiaggregant,
lodgepole pine, 221
mountain pine beetle, 221
semiochemical, 221
verbenone, 221
antibiosis,
antixenosis, 1463
crane fly, 1463
endophytes, 1463
insect behavior, 1463
antifungal activity,
Bacillus subtilis, 43
Blattella germanica, 43
identification, 43
intestinal flora, 43
antixenosis, 1463, 2602
antibiosis, 1463
Brassica oleracea, 2602
crane fly, 1463
endophytes, 1463
insect behavior, 1463
light reflectance, 2602
resistance, 2602
Thrips tabaci, 2602
aphid, 257, 1566, 1628
biorational, 257
Chromobacterium subtsugae, 257
clone dynamics, 1628
Curculio caryae, 257
development, 1628
gall, 1628
Grandevo, 257
morning glory, 1566
potyviruses, 1566
sweet potato, 1566
aphid biotype,
EPG, 2234
feeding behavior, 2234
soybean genotype, 2234
aphid damage,
economic threshold, 891
edibility, 891
lettuce crop, 891
Aphididae,
electrical penetration graph, 683
host plant suitability, 683
Miscanthus sacchariflorus, 683
phytovirus, 683
Aphidius colemani,
alphacypermethrin, 1699
cabbage, 1699
Myzus persicae, 1699
repellent netting, 1699
Aphis glycines, 1036, 1302, 1479
biotype, 1479
host-plant resistance, 1036, 1479
integrated pest management (IPM), 1302
soybean, 1479
thiamethoxam, 1302
tolerance, 1036
apical-meristem injury,
cranberry, 1339
fruit set, 1339
side-shoot, 1339
tipworm, 1339
Apis mellifera,
food availability, 2280
honey bee, 2280
hygienic behavior, 2280
seasonal effect, 2280
Apolygus lucorum,
diet, 1200
follicle, 1200
Helicoverpa armigera egg, 1200
ovarian development, 1200
apple, 520, 2118
development, 2118
diapause, 520
export, 520
life table, 2118
Panonychus ulmi, 2118
quarantine, 520
reproduction, 2118
apple cultivars,
autumn shoots, 1011
Eriosoma lanigerum (Hausmann), 1011
physiological indices, 1011
resistance to insect, 1011
apple maggot, 1173, 1511
controlled atmosphere, 1173
Diptera: Tephritidae, 1173
low oxygen, 1173
low pressure, 1173
Malus spectabilis, 1511
microsatellites, 1511
Rhagoletis indifferens, 1511
western cherry fruit fly, 1511
arthropod predators,
biological control, 1260
cotton, 1260
plant stress, 1260
whitefly, 1260

artificial diet,
Attacus atlas, 2364
hemolymph protein, 2364
Asian citrus psyllid,
Candidatus Liberibacter asiaticus, 25
citrus greening, 25
Diaphorina citri, 25
huanglongbing, 25
astigmata,
disinfestation, 1920
dispersal, 1920
integrated pest management (IPM), 1920
natural insecticide, 1920
ATP,
biochemical viability assay, 2006
colorimetric, 2006
phytosanitary treatment, 2006
quarantine, 2006
Attacus atlas,
artificial diet, 2364
hemolymph protein, 2364
Atta sexdens rubropilosa,
castor oil, 742
leaf-cutting ant, 742
physic nut, 742
seed lipid, 742
attractant compounds,
Musca domestica, 979
vinegar, 979
Auchenorrhyncha,
feeding-medium assay, 604
grapevine yellows, 604
phytoplasma-insect relationships, 604
phytoplasma seasonal variation, 604
Autographa nigrisigna,
actograph, 690
flight ability, 690
flight mill, 690
autumn shoots,
apple cultivars, 1011
Eriosoma lanigerum (Hausmann), 1011
physiological indices, 1011
resistance to insect, 1011
avocado,
phytosanitation, 2093
Raffaelea lauricola, 2093
redbay, 2093
Xyleborus glabratus, 2093
azadirachtin,
Mediterranean fruit fly, 1772
neem, 1772
South American fruit fly, 1772
toxic bait, 1772

B

Bacillus subtilis,
antifungal activity, 43
Blattella germanica, 43
identification, 43
intestinal flora, 43
Bacillus thuringiensis, 36, 1025, 1075, 1098, 1250, 1525, 2467, 2506
additives, 1075
Aedes albopictus, 1098
bioassay, 1075
biocontrol, 1075
Bt, 1025
Bt resistance, 2506
Diabrotica virgifera virgifera, 2506
dose–response assay, 2467
ELISA, 1098
field-evolved resistance, 1525
gamma radiation, 36
Helicoverpa zea, 1250
immunohistochemistry, 1098
insect resistance management, 1525, 2506
kilogray, 36
pheromone trap, 1250
Phthorimaea operculella, 1025
Plutella xylostella, 1075
receptor binding, 1098
refuge, 1525
SDS-PAGE, 36
Solanum tuberosum, 1025
toxicity, 36
toxin, 2467
transgenic, 1025
transgenic corn, 1250, 2467
western bean cutworm, 2467
Bactericera cockerelli,
Candidatus Liberibacter solanacearum, 1440
LAMP, 1440
pathogen detection, 1440
Zebra chip, 1440
Bactrocera cucurbitae,
Bactrocera dorsalis, 995
genetic sexing strain, 995
polytene chromosome, 995
Tephritidae, 995
Bactrocera dorsalis, 353, 970, 995
Bactrocera cucurbitae, 995
carboxylesterases, 970
environmental factor, 353
genetic sexing strain, 995
insecticide toxicity, 353
polytene chromosome, 995
synergist, 970
Tephritidae, 995
toxicity, 970
Bactrocera latifrons,
Capsicum annuum, 1648
cultivar, 1648
levels of infestation, 1648
Malaysian fruit fly, 1648
Bactrocera tryoni,
citrus, 883
field infestation, 883
host susceptibility, 883
bait, 929, 1794
ant, 929
boric acid, 929
Coptotermes curvignathus, 1794
Coptotermes gestroi, 1794
feeding, 929
sodium borate, 929
subterranean termite, 1794
sugar, 1794
barcode,
cockroach, 2275
invasive, 2275
Periplaneta japonica, 2275
pest, 2275
baseline susceptibility,
Chilo suppressalis, 2190
chlorantraniliprole, 2190
resistance monitoring, 2190
resistance ratio, 2190
baseline toxicity,
indoxacarb, 1423
metaflumizone, 1423
Plutella xylostella, 1423
Beauveria bassiana, 64, 1986, 2072, 2322
biological control, 64, 1986
burrower bug, 2072
chlorpyrifos, 2072
Curculionidae, 2322
Heterorhabditis bacteriophora, 2072
Listronotus maculicollis, 2322
neonicotinoid, 2322
pesticide alternative, 64, 1986
resistance management, 64, 1986
bed bug, 347, 988, 1802, 2433, 2522
carbon dioxide, 1802
chemical lure, 1802
Cimex lectularius, 347, 988
cold tolerance, 2433
control, 347
DEET, 2522
deltamethrin, 988
dust, 347
essential oil, 2522
freezing, 2433
infestation, 347
natural product, 2522
nonchemical control, 2433
pitfall, 1802
repellent, 2522
sex-specific mortality, 988
sugar, 1802
supercooling point, 2433
yeast fermentation, 1802

bee strain,
botanical origin, 1958
geographical origin, 1958
10-hydroxy-*trans*-2-decenoic acid, 1958
royal jelly, 1958
beech bark disease,
Cryptococcus fagisuga, 1324
Fagus grandifolia, 1324
invasive forest pest, 1324
beet armyworm,
insecticide resistance, 1855
resistance monitoring, 1855
behavior, 552, 2354
Bombus, 552
Diabrotica virgifera virgifera, 2354
host recognition, 2354
larva, 2354
pesticide, 552
pollinator, 552
queen-less, 552
Zea mays, 2354
Bemisia tabaci, 924, 1355, 1414, 1446, 1559
endogenous reference gene, 1446
fitness, 924
gene expression, 1446
host plant, 1414
IGRs, 1414
invasive species, 1355
neonicotinoid, 1414
nonvector herbivore, 924
population genetic, 1355
quantitative PCR, 1446
resistance, 1414
RNA-seq, 1446
tomato apex necrosis virus, 1559
tomato marchitez virus, 1559
tomato spotted wilt virus, 924
tomato torrado virus, 1559
Bemisia tabaci gossypol,
development, 1379
survival rate, 1379
bifenazate,
abamectin, 1848
acrinathrin, 1848
fenazaquin, 1848
pirimiphos methyl, 1848
bifenthrin,
Coptotermes acinaciformis, 329
Coptotermes formosanus, 329
permethrin, 329
pyrethroid, 329
binary combinations,
cyfluthrin, 1064
imidacloprid, 1064
surface treatment, 1064
binomial sequential sampling,
Acyrthosiphon pisum, 2613
dry pea, 2613
pea aphid, 2613
Pisum sativum, 2613
bioactive compound,
clove oil, 131
repellent, 131
Solenopsis invicta, 131
toxicity, 131
bioassay, 247, 1075
additives, 1075
antagonism, 247
Bacillus thuringiensis, 1075
biocontrol, 1075
horticulture, 247
pest management, 247
Plutella xylostella, 1075
synergism, 247
biochemical viability assay,
ATP, 2006
colorimetric, 2006
phytosanitary treatment, 2006
quarantine, 2006
biochemistry,
Cnaphalocrocis medinalis, 196
elemental magnesium, 196
fecundity, 196
nitrogen fertilizer, 196
biocontrol,
additives, 1075
Bacillus thuringiensis, 1075
bioassay, 1075
Plutella xylostella, 1075
bio-insecticide,
finite rate of increase, 2177
intrinsic rate of increase, 2177
net reproductive rate, 2177
population parameter, 2177
biological control, 64, 73, 80, 142, 633, 776, 1118, 1126, 1134, 1260, 1582, 1638, 1979, 1986, 1996, 2371, 2585
arthropod predators, 1260
Beauveria bassiana, 64, 1986
Bt rice, 2585
cabbage, 1134
chemical control, 142
Cleruchoides noackae, 1979
cotton, 1260
crude extract, 1118
cry gene, 2585
egg-parasitoid, 633
Eotetranychus lewisi, 80
Eulophidae, 73
fecundity, 2371
Galendromus occidentalis, 1996
generalist predator, 776, 1126
green lacewing, 1126
host age, 633
host-feeding, 1638
insect–plant interaction, 1134
integrated pest management (IPM), 1638
inundative release, 1996
kale, 1134
lethal, 776
life table, 1638, 2371
median lethal dose, 1582
median lethal time, 1582
microbial control, 1118
minimum inhibitory concentration, 1582
Mymaridae, 1979
nontarget effect, 2371
olive fruit fly, 1118
OP-replacement insecticides, 1126
Oryza sativa, 2585
parasitism, 73
parasitoid behavior, 1638
parasitoid community, 1134
pear psylla, 776
per capita reproduction, 1979
pesticide alternative, 64, 1986
Phytoseiidae, 1996
phytoseiids, 80
plant stress, 1260
Platygastridae, 73
reduced-risk insecticide, 776, 1126
resistance management, 64, 1986
secondary metabolite, 1118
strawberry, 80
sublethal effect, 776
Tetranychidae, 1996
Thaumastocoris peregrinus, 1979
total hemocyte count, 1582
toxicity risk, 142
Trichogramma spp., 142
T. urticae, 80
twospotted spider mite, 2371
Vaccinium, 73
western conifer seed bug, 633
whitefly, 1260
biological parameter,
acute toxicity, 1825
spinosad, 1825
Spodoptera exigua, 1825
sublethal effect, 1825
biophysical model,
degree-day, 124
temperature-dependent development, 124
western tarnished plant bug, 124
biorational,
aphid, 257
Chromobacterium subtsugae, 257
Curculio caryae, 257
Grandevo, 257

biosecurity,
insect, 1179
mite, 1179
phytosanitary, 1179
plant quarantine, 1179
biotic factor,
environmental factor, 2043
flight ability, 2043
flight mill, 2043
Sitodiplosis mosellana, 2043
biotype, 1479, 2595
alternative host plant, 2595
Aphis glycines, 1479
Diuraphis noxia, 2595
host-plant resistance, 1479
intrinsic rate of increase, 2595
Russian wheat aphid, 2595
soybean, 1479
bird damage,
Amyelois transitella, 1365
Corvidae, 1365
degree-day, 1365
Prunus dulcis, 1365
survivorship, 1365
blackberry,
Euschistus spp., 912
injury, 912
monitoring, 912
stink bug, 912
blacklegged tick,
granule formulation, 57
Ixodes scapularis, 57
Metarhizium brunneum, 57
microsclerotia, 57
Blatta orientalis,
blattidae, 2428
invasive species, 2428
oriental cockroach, 2428
Blattella asahinai,
harborage preference, 322
mulch, 322
toxicity, 322
Blattella germanica,
antifungal activity, 43
Bacillus subtilis, 43
identification, 43
intestinal flora, 43
blattidae,
Blatta orientalis, 2428
invasive species, 2428
oriental cockroach, 2428
Blissus occiduus,
Buchloë dactyloides, 2568
catalase, 2568
peroxidase, 2568
plant resistance, 2568
blueberry, 716, 1767
cultivar, 1767
Drosophila suzukii, 1767
Formica sp., 716
fruit firmness, 1767
insecticide toxicity, 716
mound-building ants, 716
oviposition, 1767
boll weevil,
COI DNA, 437
cotton pest, 437
DNA barcode, 437
host use, 437
Bombus,
behavior, 552
pesticide, 552
pollinator, 552
queen-less, 552
Bombus terrestris,
acquisition, 277
Metarhizium anisopliae strain F52, 277
miniature dispenser, 277
worker mortality, 277
booklouse,
insect pest, 491
Liposcelis, 491
Psocoptera, 491
stored product, 491
bootstrap,
Chrysopa pallens, 1
jackknife, 1
life table, 1
population projection, 1
bootstrap percentile,
lower developmental threshold, 107
Performance-2 model, 107
sum of effective temperatures, 107
upper developmental threshold, 107
borer damage,
fumonisin, 2241
Fusarium verticillioides, 2241
Sesamia nonagrioides, 2241
Zea mays, 2241
boric acid,
ant, 929
bait, 929
feeding, 929
sodium borate, 929
boring larva,
branch protection zone, 1602
natural target pruning, 1602
olive pyralid moth, 1602
pruning paint, 1602
boscalid,
almond, 1952
diflubenzuron, 1952
organosilicone, 1952
pyraclostrobin, 1952
botanical insecticides,
essential oil, 159
GC-MS, 159
scarab larvae, 159
white grub, 159
botanical origin,
bee strain, 1958
geographical origin, 1958
10-hydroxy-*trans*-2-decenoic acid, 1958
royal jelly, 1958
bottle trap,
ambrosia beetle, 1780
forecasting activity, 1780
phenological indicator, 1780
trap tree, 1780
Brachiaria,
adult damage, 1871
host-plant resistance, 1871
spittlebug, 1871
branch protection zone,
boring larva, 1602
natural target pruning, 1602
olive pyralid moth, 1602
pruning paint, 1602
Brassica oleracea,
antixenosis, 2602
light reflectance, 2602
resistance, 2602
Thrips tabaci, 2602
breeding system,
invasive species, 2216
neoteny, 2216
termite, 2216
Brevicoryne brassicae,
insecticide resistance, 954
neonicotinoid, 954
organophosphate, 954
pyrethroid, 954
broccoli, 267, 1103, 1726
cabbage, 267
cauliflower, 1726
Diadegma insulare, 1103
diamondback moth, 1103
diel activity, 1726
insecticide timing, 267
integrated pest management (IPM), 267
invasive species, 267
pest management, 1726
Plutella xylostella, 1103
simulation, 1103
stink bug, 1726
bronze beetle,
Chrysomelidae, 899
degree-day, 899
phenology model, 899
temperature, 899
brown marmorated stink bug, 150, 2141
invasive species, 150
maximum temperature, 2141
movement, 150
overwintering adult, 2141
pesticide, 150
population density, 2141
sublethal, 150

Bt,
Bacillus thuringiensis, 1025
Phthorimaea operculella, 1025
Solanum tuberosum, 1025
transgenic, 1025
Bt cotton, 375, 1819
emamectin benzoate, 375
fenvalerate, 375
Helicoverpa armigera, 375
insect resistance management, 1819
phoxim, 375
tobacco budworm, 1819
Bt plant,
genetic engineering, 2151
insecticide reduction, 2151
Lepidoptera, 2151
maize, 2151
Bt resistance,
Bacillus thuringiensis, 2506
Diabrotica virgifera virgifera, 2506
insect resistance management, 2506
Bt rice, 1887, 2585
biological control, 2585
cry gene, 2585
detoxification enzyme, 1887
ecological fitness, 1887
Nilaparvata lugens, 1887
Oryza sativa, 2585
population, 1887
Bt toxin,
Helicoverpa armigera, 614
lifespan, 614
reproductive performance, 614
Spodoptera exigua, 614
Buchloë dactyloides,
Blissus occiduus, 2568
catalase, 2568
peroxidase, 2568
plant resistance, 2568
bullet gall,
alternating generation, 1747
gall wasp, 1747
Quercus virginiana, 1747
burrower bug,
Beauveria bassiana, 2072
chlorpyrifos, 2072
Heterorhabditis bacteriophora, 2072

C

cabbage, 267, 1134, 1699
alphacypermethrin, 1699
Aphidius colemani, 1699
biological control, 1134
broccoli, 267
insecticide timing, 267
insect–plant interaction, 1134
integrated pest management (IPM), 267
invasive species, 267
kale, 1134
Myzus persicae, 1699
parasitoid community, 1134
repellent netting, 1699
Cacopsylla chinensis, 536, 1349
Allium sativum, 1349
allyl trisulfide, 1349
contact toxicity, 1349
cytochrome oxidase I, 536
diallyl disulfide, 1349
16S rDNA, 536
Callosobruchus maculatus,
cowpea, 2473
insect resistance management, 2473
simulation, 2473
transgenic crop, 2473
Candidatus Liberibacter asiaticus,
Asian citrus psyllid, 25
citrus greening, 25
Diaphorina citri, 25
huanglongbing, 25
Candidatus Liberibacter solanacearum,
Bactericera cockerelli, 1440
LAMP, 1440
pathogen detection, 1440
Zebra chip, 1440
canopy translocation,
hemlock woolly adelgid, 2399
neonicotinoid, 2399
systemic insecticide, 2399
Tsuga canadensis, 2399
Capsicum annuum,
Bactrocera latifrons, 1648
cultivar, 1648
levels of infestation, 1648
Malaysian fruit fly, 1648
carbamate,
Myzus persicae, 1838
oilseed rape, 1838
pyrethroid, 1838
target site resistance, 1838
carbon dioxide,
bed bug, 1802
chemical lure, 1802
pitfall, 1802
sugar, 1802
yeast fermentation, 1802
carboxylesterases,
Bactrocera dorsalis, 970
synergist, 970
toxicity, 970
carcass-burying behavior,
Coptotermes formosanus, 1812
disease resistance, 1812
lufenuron, 1812
Pseudomonas aeruginosa, 1812
carrion,
forensic entomology, 1927
myiasis, 1927
secondary screwworm, 1927
spray-dried blood, 1927
castor oil,
Atta sexdens rubropilosa, 742
leaf-cutting ant, 742
physic nut, 742
seed lipid, 742
catalase,
Blissus occiduus, 2568
Buchloë dactyloides, 2568
peroxidase, 2568
plant resistance, 2568
cauliflower,
broccoli, 1726
diel activity, 1726
pest management, 1726
stink bug, 1726
centroid size,
crop rotation resistance, 1517
Diabrotica virgifera virgifera, 1517
dispersal, 1517
Cerambycidae,
Cleridae, 206
Histeridae, 206
Scolytinae, 206
Trogossitidae, 206
cereal crops,
cultural control, 1294
English grain aphid, 1294
host switch, 1294
life table, 1294
Cheilomenes sexmaculata development,
adult longevity, 2621
fertility, 2621
prey consumption, 2621
chemical control, 142, 2286
biological control, 142
exotic insect, 2286
insect–disease complex, 2286
Lauraceae, 2286
redbay ambrosia beetle, 2286
toxicity risk, 142
Trichogramma spp., 142
chemical lure,
bed bug, 1802
carbon dioxide, 1802
pitfall, 1802
sugar, 1802
yeast fermentation, 1802
Chilo suppressalis, 1832, 2190
baseline susceptibility, 2190
chlorantraniliprole, 2190
dynamics, 1832
insecticide resistance, 1832
regression analysis, 1832
resistance monitoring, 2190
resistance ratio, 2190

chitinase inhibitor,
eastern subterranean termite, 1788
toxicity, 1788
chlorantraniliprole, 181, 2190
baseline susceptibility, 2190
Chilo suppressalis, 2190
resistance monitoring, 2190
resistance ratio, 2190
rice water weevil, 181
seed treatment, 181
sublethal effect, 181
thiamethoxam, 181
chlorpyrifos, 1226, 2072
Beauveria bassiana, 2072
burrower bug, 2072
Heterorhabditis bacteriophora, 2072
pesticide toxicity, 1226
spinosad, 1226
sterile insect technique, 1226
Vienna-8, 1226
chlorpyrifos-methyl plus deltamethrin,
field strains, 1911
mortality, 1911
spinosad, 1911
stored-grain insects, 1911
choice test,
cole crop, 283
harlequin bug, 283
Murgantia histrionica, 283
trap crop, 283
Chromobacterium subtsugae,
aphid, 257
biorational, 257
Curculio caryae, 257
Grandevo, 257
Chrysomelidae,
bronze beetle, 899
degree-day, 899
phenology model, 899
temperature, 899
Chrysopa pallens,
bootstrap, 1
jackknife, 1
life table, 1
population projection, 1
Cicadellidae,
Dracaena marginata, 2027
nursery stock, 2027
offshore preclearance program, 2027
size restriction, 2027
Cimex lectularius, 347, 988
bed bug, 347, 988
control, 347
deltamethrin, 988
dust, 347
infestation, 347
sex-specific mortality, 988
circadian clock,
clock genes, 450
cryptochrome, 450
Helicoverpa armigera, 450
photoreceptor, 450
citrus,
Bactrocera tryoni, 883
field infestation, 883
host susceptibility, 883
citrus greening,
Asian citrus psyllid, 25
Candidatus Liberibacter asiaticus, 25
Diaphorina citri, 25
huanglongbing, 25
Cleridae,
Cerambycidae, 206
Histeridae, 206
Scolytinae, 206
Trogossitidae, 206
Cleruchoides noackae,
biological control, 1979
Mymaridae, 1979
per capita reproduction, 1979
Thaumastocoris peregrinus, 1979
clock genes,
circadian clock, 450
cryptochrome, 450
Helicoverpa armigera, 450
photoreceptor, 450
clone dynamics,
aphid, 1628
development, 1628
gall, 1628
clove oil,
bioactive compound, 131
repellent, 131
Solenopsis invicta, 131
toxicity, 131
clover,
Coleophora deauratella, 844
monitoring, 844
pheromone-baited traps, 844
Cnaphalocrocis medinalis,
biochemistry, 196
elemental magnesium, 196
fecundity, 196
nitrogen fertilizer, 196
cockroach,
barcode, 2275
invasive, 2275
Periplaneta japonica, 2275
pest, 2275
codling moth,
diagnostic concentration, 939
insecticide resistance, 939
neonate, 939
coffee berry borer,
freezing, 653
green coffee, 653
Hawaii, 653
Hypothenemus hampei, 653
COI,
genetic distance, 408
phylogenetic analysis, 408
thrip, 408
COI DNA,
boll weevil, 437
cotton pest, 437
DNA barcode, 437
host use, 437
cold tolerance,
bed bug, 2433
freezing, 2433
nonchemical control, 2433
supercooling point, 2433
cold treatment, 1608, 2336
Mediterranean fruit fly, 2336
peach fruit fly, 1608, 2336
phytosanitary treatment, 1608, 2336
phytosanitation, 1608
quarantine treatment, 1608, 2336
cole crop,
choice test, 283
harlequin bug, 283
Murgantia histrionica, 283
trap crop, 283
Coleophora deauratella,
clover, 844
monitoring, 844
pheromone-baited traps, 844
Colorado potato beetle,
diapause, 2491
insecticide resistance, 2491
integrated pest management (IPM), 2491
phenology, 2491
colorimetric,
ATP, 2006
biochemical viability assay, 2006
phytosanitary treatment, 2006
quarantine, 2006
compensation,
corn, 816
Diatraea grandiosella, 816
refuge, 816
southwestern corn borer, 816
cone abortion,
Ernobius spp., 229
Leptoglossus occidentalis, 229
nut production, 229
stone pine, 229
conservation biological control,
flowering plant, 2327
habitat management, 2327
natural enemy, 2327
Sphaerophoria spp., 2327
Conservation Reserve Program,
Alaska, 669
grasshopper bait, 669
high-latitude agriculture, 669
Insecta, 669

contact toxicity,
Allium sativum, 1349
allyl trisulfide, 1349
Cacopsylla chinensis, 1349
diallyl disulfide, 1349
control, 347, 1545
ant, 1545
bed bug, 347
Cimex lectularius, 347
dust, 347
eradication, 1545
infestation, 347
invasive ants, 1545
treatment, 1545
controlled atmosphere,
apple maggot, 1173
Diptera: Tephritidae, 1173
low oxygen, 1173
low pressure, 1173
Coptotermes,
nylon 12, 1395
plastic cable sheathings, 1395
polyethylene, 1395
subterranean termite, 1395
Coptotermes acinaciformis,
bifenthrin, 329
Coptotermes formosanus, 329
permethrin, 329
pyrethroid, 329
Coptotermes curvignathus,
bait, 1794
Coptotermes gestroi, 1794
subterranean termite, 1794
sugar, 1794
Coptotermes formosanus, 329, 1812, 1863, 2442, 2630
bifenthrin, 329
carcass-burying behavior, 1812
Coptotermes acinaciformis, 329
differentially expressed gene, 1863
disease resistance, 1812
fipronil, 2630
group size, 2630
horizontal transfer, 2630
insecticide, 1863
Isoptera, 2442
lufenuron, 1812
mark–recapture protocol, 2442
permethrin, 329
probability density function, 2442
Pseudomonas aeruginosa, 1812
pyrethroid, 329
Rhinotermitidae, 2442
Coptotermes gestroi,
bait, 1794
Coptotermes curvignathus, 1794
subterranean termite, 1794
sugar, 1794
corn, 816, 2048
compensation, 816
Diatraea grandiosella, 816
Japanese beetle, 2048
Popillia japonica, 2048
refuge, 816
silk clipping, 2048
southwestern corn borer, 816
corn leafhopper,
incidence, 1574
mollicutes, 1574
overwintering, 1574
pest, 1574
cortical parenchyma,
Phloeomyzus passerinii, 473
Populus x canadensis, 473
tree resistance, 473
Corvidae,
Amyelois transitella, 1365
bird damage, 1365
degree-day, 1365
Prunus dulcis, 1365
survivorship, 1365
cotton, 86, 189, 862, 1209, 1260, 1286, 2378
arthropod predators, 1260
biological control, 1260
Creontiades signatus, 189
cultural control, 2378
ecoinformatics, 1286
farmer decision-making, 1286
feeding injury, 189
insect damage, 1209
lint yield, 1209
Lygus, 1286
Lygus hesperus, 1209
Lygus lineolaris, 2378
nonreproducing state, 86
observational studies, 1286
plant compensation, 1209
plant stress, 1260
Tetranychus urticae, 862
tropical population, 86
twospotted spider mite, 862
verde plant bug, 189
whitefly, 1260
yield, 862
cotton aphid,
flonicamid, 1430
neonicotinoid, 1430
resistance, 1430
sulfoxaflor, 1430
cotton boll damage,
pest monitoring, 1310
plant bug, 1310
sucking bugs, 1310
cotton boll injury,
Gossypium hirsutum, 2077
insect–plant interaction, 2077
plant bug, 2077
cotton fleahopper,
Gossypium hirsutum, 1045
plant resistance, 1045
Pseudatomoscelis seriatus, 1045
cotton pest, 437, 2225
AFLP, 2225
boll weevil, 437
COI DNA, 437
dispersal, 2225
DNA barcode, 437
gene flow, 2225
host use, 437
population structure, 2225
courtship,
mating behavioral sequence, 1373
mating success, 1373
sexual selection, 1373
cowpea,
Callosobruchus maculatus, 2473
insect resistance management, 2473
simulation, 2473
transgenic crop, 2473
Crambe spp.,
flea beetle, 2557
glucosinolate, 2557
Phyllotreta spp., 2557
Sinapis alba, 2557
cranberry, 1339, 2424
apical-meristem injury, 1339
fruit, 2424
fruit set, 1339
ripeness, 2424
side-shoot, 1339
spotted wing drosophila, 2424
susceptibility, 2424
tipworm, 1339
crane fly,
antibiosis, 1463
antixenosis, 1463
endophytes, 1463
insect behavior, 1463
Creontiades signatus,
cotton, 189
feeding injury, 189
verde plant bug, 189
crop pollination,
agricultural landscape, 97
seminatural areas pollinator habitat, 97
wild bees, 97
crop rotation resistance,
centroid size, 1517
Diabrotica virgifera virgifera, 1517
dispersal, 1517
cross-resistance, 959, 1029, 2167
Aedes aegypti, 959
dieldrin, 2167
fipronil, 2167
insecticide resistance, 959, 2167

native species, 1029
plant resistance, 1029
pyrethroid, 959
Rdl gene, 2167
wheat pests, 1029
crude extract,
biological control, 1118
microbial control, 1118
olive fruit fly, 1118
secondary metabolite, 1118
cry gene,
biological control, 2585
Bt rice, 2585
Oryza sativa, 2585
Cry1Ab/Cry2Aj protein,
Cry1Ab/Vip3DA protein, 2548
insect resistance evaluation, 2548
Ostrinia furnacalis, 2548
transgenic maize, 2548
Cry1Ab/Vip3DA protein,
Cry1Ab/Cry2Aj protein, 2548
insect resistance evaluation, 2548
Ostrinia furnacalis, 2548
transgenic maize, 2548
Cry1Ac resistance,
Helicoverpa armigera, 2514
nonbinding site, 2514
proteinases, 2514
cryopreservation,
embryo, 855
Pectinophora gossypiella, 855
permeabilization, 855
pink bollworm moth, 855
cryptic species,
cytochrome *c* oxidase subunit I, 463
DNA-based identification, 463
internal transcribed spacer, 463
Tetranychus, 463
cryptochrome,
circadian clock, 450
clock genes, 450
Helicoverpa armigera, 450
photoreceptor, 450
Cryptococcus fagisuga,
beech bark disease, 1324
Fagus grandifolia, 1324
invasive forest pest, 1324
CT scan,
damage progression, 1331
fire-killed trees, 1331
salvage logging, 1331
whitespotted sawyer, 1331
Culex pipiens pallens,
knockdown resistance, 360
pyrethroid insecticide resistance, 360
susceptibility, 360
cultivar, 1648, 1767
Bactrocera latifrons, 1648
blueberry, 1767
Capsicum annuum, 1648
Drosophila suzukii, 1767
fruit firmness, 1767
levels of infestation, 1648
Malaysian fruit fly, 1648
oviposition, 1767
cultural control, 1294, 2378
cereal crops, 1294
cotton, 2378
English grain aphid, 1294
host switch, 1294
life table, 1294
Lygus lineolaris, 2378
Curculio caryae,
aphid, 257
biorational, 257
Chromobacterium subtsugae, 257
Grandevo, 257
Curculionidae,
Beauveria bassiana, 2322
Listronotus maculicollis, 2322
neonicotinoid, 2322
cutworm,
Abagrotis, 2135
grape pest management, 2135
insecticide toxicity, 2135
cyantraniliprole,
Cyazypyr, 597
fluorescein sodium salt, 597
imidacloprid, 597
spirotetramat, 597
Cyazypyr,
cyantraniliprole, 597
fluorescein sodium salt, 597
imidacloprid, 597
spirotetramat, 597
cyfluthrin, 400, 1064
binary combinations, 1064
imidacloprid, 400, 1064
metamidophos, 400
Myzus persicae, 400
pirimicarb, 400
surface treatment, 1064
Cylas formicarius,
Diabrotica, 1456
Ipomoea, 1456
plant resistance, 1456
white grub, 1456
CYP4,
cytochrome P450 monooxygenase, 393
Diaphorina citri, 393
gene expression, 393
protein expression, 393
cytochrome *c* oxidase subunit I,
cryptic species, 463
DNA-based identification, 463
internal transcribed spacer, 463
Tetranychus, 463
cytochrome oxidase I,
Cacopsylla chinensis, 536
16S rDNA, 536
cytochrome P450 monooxygenase,
CYP4, 393
Diaphorina citri, 393
gene expression, 393
protein expression, 393

D

daily reproductive rhythm,
adult starvation, 215
Glenea cantor, 215
lifetime reproductive pattern, 215
sexual maturation, 215
damage,
mortality, 786
potato, 786
recovery, 786
wireworm, 786
damage progression,
CT scan, 1331
fire-killed trees, 1331
salvage logging, 1331
whitespotted sawyer, 1331
DAMs,
flight initiation, 2259
insecticide resistance, 2259
walking, 2259
dead-heart plants,
Diabrotica undecimpunctata howardi, 1669
Maize, 1669
seedling damage, 1669
southern corn rootworm, 1669
debarking,
firewood, 235
grinding, 235
oak mortality, 235
solarization, 235
DEET,
bed bug, 2522
essential oil, 2522
natural product, 2522
repellent, 2522
defoliation,
Glycine max, 830
insect resistance, 830
value added traits, 830
degree-day, 124, 899, 1365, 1495, 1757
action threshold, 1757
Amyelois transitella, 1365
biophysical model, 124
bird damage, 1365
bronze beetle, 899
Chrysomelidae, 899

Corvidae, 1365
distribution, 1495
Fragaria ananassa, 1757
invasive species ecology, 1495
mite-day, 1757
phenology model, 899
Prunus dulcis, 1365
six-spotted thrip, 1757
survey, 1495
survivorship, 1365
temperature, 899
temperature-dependent development, 124
western tarnished plant bug, 124
deltamethrin,
bed bug, 988
Cimex lectularius, 988
sex-specific mortality, 988
demographic parameter,
actuarial mathematics, 675
Anarsia lineatella, 675
intrinsic rate of increase, 675
life table, 675
desert environment,
host plant, 1404
insecticide resistance, 1404
systemic uptake bioassay, 1404
detection capacity,
environmental risk assessment, 1659
insect, 1659
statistical power, 1659
transgenic crop, 1659
detection tools,
multifunnel trap, 2407
prism trap, 2407
Sabic green, 2407
Sabic purple, 2407
detoxification enzyme,
Bt rice, 1887
ecological fitness, 1887
Nilaparvata lugens, 1887
population, 1887
development, 1379, 1628, 2101, 2118
Agrilus planipennis, 2101
aphid, 1628
apple, 2118
Bemisia tabaci gossypol, 1379
clone dynamics, 1628
Fraxinus uhdei, 2101
gall, 1628
invasive, 2101
laboratory rearing, 2101
life table, 2118
Panonychus ulmi, 2118
reproduction, 2118
survival rate, 1379
Diabrotica,
Cylas formicarius, 1456
Ipomoea, 1456
plant resistance, 1456
white grub, 1456
Diabrotica barberi, 168, 622, 1941
Diabrotica virgifera virgifera, 168, 622, 1941
economic injury level, 168
geographic information systems, 168
Heterorhabditis, 622
integrated pest management (IPM), 168
Metarhizium, 622
pesticide mixture, 1941
pyramid, 1941
resistance management, 1941
Steinernema, 622
Diabrotica undecimpunctata howardi,
dead-heart plants, 1669
Maize, 1669
seedling damage, 1669
southern corn rootworm, 1669
Diabrotica virgifera virgifera, 168, 622, 823, 1517, 1941, 2195, 2354, 2506
Bacillus thuringiensis, 2506
behavior, 2354
Bt resistance, 2506
centroid size, 1517
crop rotation resistance, 1517
Diabrotica barberi, 168, 622, 1941
dispersal, 1517
economic injury level, 168
event 59122, 2195
Event MON 88017, 823
Event MON 89034, 823
geographic information systems, 168
Heterorhabditis, 622
host recognition, 2354
insect resistance management, 2506
integrated pest management (IPM), 168
larva, 2354
Metarhizium, 622
pesticide mixture, 1941
pyramid, 1941
refuge strategy, 2195
resistance management, 823, 1941, 2195
Steinernema, 622
transgenic corn, 823
Zea mays, 2354
Diadegma insulare,
broccoli, 1103
diamondback moth, 1103
Plutella xylostella, *1103*
simulation, 1103
diagnostic concentration,
codling moth, 939
insecticide resistance, 939
neonate, 939
diallyl disulfide,
Allium sativum, 1349
allyl trisulfide, 1349
Cacopsylla chinensis, 1349
contact toxicity, 1349
diamondback moth,
broccoli, 1103
Diadegma insulare, 1103
Plutella xylostella, *1103*
simulation, 1103
diapause, 520, 1089, 2491
apple, 520
Colorado potato beetle, 2491
export, 520
insecticide resistance, 2491
integrated pest management (IPM), 2491
Megachile rotundata, 1089
metabolic rate, 1089
phenology, 2491
quarantine, 520
respiration, 1089
thermal stress, 1089
Diaphorina citri, 25, 393
Asian citrus psyllid, 25
Candidatus Liberibacter asiaticus, 25
citrus greening, 25
CYP4, 393
cytochrome P450 monooxygenase, 393
gene expression, 393
huanglongbing, 25
protein expression, 393
Diatraea grandiosella,
compensation, 816
corn, 816
refuge, 816
southwestern corn borer, 816
diel activity,
broccoli, 1726
cauliflower, 1726
pest management, 1726
stink bug, 1726
dieldrin,
cross-resistance, 2167
fipronil, 2167
insecticide resistance, 2167
Rdl gene, 2167
diet,
Apolygus lucorum, 1200
follicle, 1200
Helicoverpa armigera egg, 1200
ovarian development, 1200
differentially expressed gene,
Coptotermes formosanus, 1863
insecticide, 1863

diflubenzuron,
almond, 1952
boscalid, 1952
organosilicone, 1952
pyraclostrobin, 1952
digestive enzymes,
nutritional index, 1018
Phthorimaea operculella, 1018
potato germplasms, 1018
Diglyphus isaea,
greenhouse biocontrol, 1590
greenhouse pest management, 1590
leafminer biocontrol, 1590
safe pesticide, 1590
Diptera: Tephritidae,
apple maggot, 1173
controlled atmosphere, 1173
low oxygen, 1173
low pressure, 1173
disease resistance,
carcass-burying behavior, 1812
Coptotermes formosanus, 1812
lufenuron, 1812
Pseudomonas aeruginosa, 1812
disinfestation,
astigmata, 1920
dispersal, 1920
integrated pest management (IPM), 1920
natural insecticide, 1920
dispersal, 1517, 1920, 2225, 2296
AFLP, 2225
astigmata, 1920
centroid size, 1517
cotton pest, 2225
crop rotation resistance, 1517
Diabrotica virgifera virgifera, 1517
disinfestation, 1920
gene flow, 2225
Hemiptera vector, 2296
integrated pest management (IPM), 1920
natural insecticide, 1920
plant community, 2296
population structure, 2225
vineyard agroecosystem, 2296
distribution, 874, 1495, 2084
Acer platanoides, 874
degree-day, 1495
imidacloprid, 874
invasive species ecology, 1495
neonicotinoid, 874
Ostrinia nubilalis, 2084
pheromone, 2084
predictive model, 2084
spatial, 2084
survey, 1495
translocation, 874
Diuraphis noxia,
alternative host plant, 2595
biotype, 2595
intrinsic rate of increase, 2595
Russian wheat aphid, 2595
DNA barcode,
boll weevil, 437
COI DNA, 437
cotton pest, 437
host use, 437
DNA-based identification,
cryptic species, 463
cytochrome *c* oxidase subunit I, 463
internal transcribed spacer, 463
Tetranychus, 463
dose–response,
electroantennogram, 1653
floral compounds, 1653
linalool, 1653
dose–response assay,
Bacillus thuringiensis, 2467
toxin, 2467
transgenic corn, 2467
western bean cutworm, 2467
Dracaena marginata,
Cicadellidae, 2027
nursery stock, 2027
offshore preclearance program, 2027
size restriction, 2027
Drosophila suzukii,
blueberry, 1767
cultivar, 1767
fruit firmness, 1767
oviposition, 1767
dry pea,
Acyrthosiphon pisum, 2613
binomial sequential sampling, 2613
pea aphid, 2613
Pisum sativum, 2613
drywood termites,
acoustic emissions, 338
feeding periodicity, 338
temperature effects, 338
dust,
bed bug, 347
Cimex lectularius, 347
control, 347
infestation, 347
dynamics,
Chilo suppressalis, 1832
insecticide resistance, 1832
regression analysis, 1832

E

eastern subterranean termite,
chitinase inhibitor, 1788
toxicity, 1788
eastern white pine,
Matsucoccidae, 2391
Matsucoccus macrocicatrices, 2391
Pinus strobus, 2391
southeastern United States, 2391
Echinacea angustifolia,
insect pollinator, 2055
pollen germination, 2055
pollen-tube quantification, 2055
shrivelled (retracted) styles, 2055
ecoinformatics,
cotton, 1286
farmer decision-making, 1286
Lygus, 1286
observational studies, 1286
ecological fitness,
Bt rice, 1887
detoxification enzyme, 1887
Nilaparvata lugens, 1887
population, 1887
economic,
fitness costs, 366
insecticide resistance, 366
malaria, 366
vector control, 366
economic analysis,
esfenvalerate, 1503
methoprene, 1503
Plodia interpunctella, 1503
economic injury level, 168, 905, 1274
Diabrotica barberi, 168
Diabrotica virgifera virgifera, 168
economic threshold, 1274
geographic information systems, 168
grape, 905
integrated pest management (IPM), 168, 905
larval survival, 1274
Paralobesia viteana, 905
Striacosta albicosta, 1274
western bean cutworm, 1274
economic threshold, 891, 1274
aphid damage, 891
economic injury level, 1274
edibility, 891
larval survival, 1274
lettuce crop, 891
Striacosta albicosta, 1274
western bean cutworm, 1274
edibility,
aphid damage, 891
economic threshold, 891
lettuce crop, 891

egg,
embryology, 747
fall armyworm, 747
insecticide, 747
morphology, 747
egg parasitism,
Gryon japonicum, 1973
kairomone, 1973
Ooencyrtus nezarae, 1973
soybean, 1973
egg-parasitoid,
biological control, 633
host age, 633
western conifer seed bug, 633
electrical penetration graph,
Aphididae, 683
host plant suitability, 683
Miscanthus sacchariflorus, 683
phytovirus, 683
electroantennogram,
dose–response, 1653
floral compounds, 1653
linalool, 1653
electronic penetration graph,
seed treatment, 2384
soybean aphid, 2384
thiamethoxam, 2384
elemental magnesium,
biochemistry, 196
Cnaphalocrocis medinalis, 196
fecundity, 196
nitrogen fertilizer, 196
ELISA,
Aedes albopictus, 1098
Bacillus thuringiensis, 1098
immunohistochemistry, 1098
receptor binding, 1098
emamectin benzoate,
Bt cotton, 375
fenvalerate, 375
Helicoverpa armigera, 375
phoxim, 375
embryo,
cryopreservation, 855
Pectinophora gossypiella, 855
permeabilization, 855
pink bollworm moth, 855
embryology,
egg, 747
fall armyworm, 747
insecticide, 747
morphology, 747
Endoclita signifier,
Eucalyptus, 866
forest pest, 866
life cycle, 866
morphology, 866
endogenous reference gene,
Bemisia tabaci, 1446
gene expression, 1446
quantitative PCR, 1446
RNA-seq, 1446
endophytes,
antibiosis, 1463
antixenosis, 1463
crane fly, 1463
insect behavior, 1463
English grain aphid, 1294, 1473
cereal crops, 1294
cultural control, 1294
germplasm, 1473
host switch, 1294
life table, 1294
resistance, 1473
tolerance, 1473
wheat, 1473
entomopathogenic nematode, 648, 1112
Heterorhabditis, 1112
integrated pest management (IPM), 648
peanut, 1112
Steinernema, 1112
Steinernema carpocapsae, 648
Steinernematidae, 648
Tenebrionidae, 648
white grub, 1112
environmental factor, 353, 2043
Bactrocera dorsalis (Hendel), 353
biotic factor, 2043
flight ability, 2043
flight mill, 2043
insecticide toxicity, 353
Sitodiplosis mosellana, 2043
environmental risk assessment,
detection capacity, 1659
insect, 1659
statistical power, 1659
transgenic crop, 1659
Eotetranychus lewisi,
biological control, 80
phytoseiids, 80
strawberry, 80
T. urticae, 80
EPG,
aphid biotype, 2234
feeding behavior, 2234
soybean genotype, 2234
Epimedium pubescens,
Liposcelis bostrychophila, 513
(R)-carvone, 513
repellency, 513
Tribolium castaneum, 513
Epiphyas postvittana,
methyl bromide, 1155
postharvest fumigation, 1155
sorption kinetics, 1155
eradication,
ant, 1545
control, 1545
invasive ants, 1545
treatment, 1545
Eriosoma lanigerum (Hausmann),
apple cultivars, 1011
autumn shoots, 1011
physiological indices, 1011
resistance to insect, 1011
Ernobius spp.,
cone abortion, 229
Leptoglossus occidentalis, 229
nut production, 229
stone pine, 229
Erotylidae,
Europs, 305
Loberus testaceus, 305
Nitidulidae, 305
Omatomidae, 305
esfenvalerate,
economic analysis, 1503
methoprene, 1503
Plodia interpunctella, 1503
essential oil, 159, 1902, 2522
bed bug, 2522
botanical insecticides, 159
DEET, 2522
GC-MS, 159
Juniperus oxycedrus ssp. *oxycedrus*, 1902
natural product, 2522
repellent, 2522
scarab larvae, 159
silica gel, 1902
Sitophilus oryzae, 1902
stored wheat, 1902
Tribolium confusum, 1902
white grub, 159
establishment,
invasive, 1145
natural enemy introduction, 1145
parasitoid release, 1145
wood borers, 1145
ethanol,
α-pinene, 1684
ipsdienol, 1684
ipsenol, 1684
kairomone, 1684
ethyl acetate,
integrated pest management (IPM), 1739
kairomone, 1739
red palm weevil, 1739
trap, 1739
Eucalyptus,
Endoclita signifier, 866
forest pest, 866
life cycle, 866
morphology, 866
Eulophidae,
biological control, 73
parasitism, 73
Platygastridae, 73
Vaccinium, 73

European corn borer,
maize, 1234
Mediterranean corn borer, 1234
plant development, 1234
resistance, 1234
Europs,
Erotylidae, 305
Loberus testaceus, 305
Nitidulidae, 305
Omatomidae, 305
Euschistus spp.,
blackberry, 912
injury, 912
monitoring, 912
stink bug, 912
event 59122,
Diabrotica virgifera virgifera, 2195
refuge strategy, 2195
resistance management, 2195
Event MON 88017,
Diabrotica virgifera virgifera, 823
Event MON 89034, 823
resistance management, 823
transgenic corn, 823
Event MON 89034,
Diabrotica virgifera virgifera, 823
Event MON 88017, 823
resistance management, 823
transgenic corn, 823
exotic insect,
chemical control, 2286
insect–disease complex, 2286
Lauraceae, 2286
redbay ambrosia beetle, 2286
export,
apple, 520
diapause, 520
quarantine, 520

F

F1538I mutation,
fenpropathrin, 2457
molecular marker, 2457
Tetranychus cinnabarinus, 2457
voltage-gate sodium channel, 2457
Fagus grandifolia,
beech bark disease, 1324
Cryptococcus fagisuga, 1324
invasive forest pest, 1324
fall armyworm,
egg, 747
embryology, 747
insecticide, 747
morphology, 747
farmer decision-making,
cotton, 1286
ecoinformatics, 1286
Lygus, 1286
observational studies, 1286
fat body,
honey bee, 1553
nutritional status, 1553
pollen diets, 1553
protein, 1553
fecundity, 10, 196, 2371
biochemistry, 196
biological control, 2371
Cnaphalocrocis medinalis, 196
elemental magnesium, 196
life table, 2371
Nilaparvata lugens, 10
nitrogen fertilizer, 196
nontarget effect, 2371
sublethal effect, 10
survival, 10
twospotted spider mite, 2371
wing formation, 10
feeding,
ant, 929
bait, 929
boric acid, 929
sodium borate, 929
feeding behavior,
aphid biotype, 2234
EPG, 2234
soybean genotype, 2234
feeding guild,
injury equivalency, 837
Oryza sativa, 837
rice, 837
rice planthopper, 837
feeding injury,
cotton, 189
Creontiades signatus, 189
verde plant bug, 189
feeding-medium assay,
Auchenorrhyncha, 604
grapevine yellows, 604
phytoplasma-insect relationships, 604
phytoplasma seasonal variation, 604
feeding periodicity,
acoustic emissions, 338
drywood termites, 338
temperature effects, 338
fenazaquin,
abamectin, 1848
acrinathrin, 1848
bifenazate, 1848
pirimiphos methyl, 1848
fenpropathrin,
F1538I mutation, 2457
molecular marker, 2457
Tetranychus cinnabarinus, 2457
voltage-gate sodium channel, 2457
fenvalerate,
Bt cotton, 375
emamectin benzoate, 375
Helicoverpa armigera, 375
phoxim, 375
Ferrisia gilli,
Acerophagus, 1716
grape, 1716
mealybug, 1716
Vitis, 1716
fertility,
adult longevity, 2621
Cheilomenes sexmaculata development, 2621
prey consumption, 2621
field-evolved resistance,
Bacillus thuringiensis, 1525
insect resistance management, 1525
refuge, 1525
field infestation,
Bactrocera tryoni, 883
citrus, 883
host susceptibility, 883
field strains,
chlorpyrifos-methyl plus deltamethrin, 1911
mortality, 1911
spinosad, 1911
stored-grain insects, 1911
finite rate of increase,
bio-insecticide, 2177
intrinsic rate of increase, 2177
net reproductive rate, 2177
population parameter, 2177
fipronil, 2160, 2167, 2630
Coptotermes formosanus, 2630
cross-resistance, 2167
dieldrin, 2167
foraging behavior, 2160
group size, 2630
horizontal transfer, 2630
insecticide resistance, 2167
Rdl gene, 2167
Reticulitermes, 2160
termite, 2160
termiticide, 2160
fire-killed trees,
CT scan, 1331
damage progression, 1331
salvage logging, 1331
whitespotted sawyer, 1331
firewood,
debarking, 235
grinding, 235
oak mortality, 235
solarization, 235
first-generation,
life table, 533
Megacopta cribraria, 533
rearing, 533
soybean, 533

fitness, 924, 2310
Bemisia tabaci, 924
generalist predator, 2310
inundative release, 2310
invasive species, 2310
mass-rearing, 2310
nonvector herbivore, 924
tomato spotted wilt virus, 924
fitness costs,
economic, 366
insecticide resistance, 366
malaria, 366
vector control, 366
flea beetle,
Crambe spp., 2557
glucosinolate, 2557
Phyllotreta spp., 2557
Sinapis alba, 2557
flight ability, 690, 2043
actograph, 690
Autographa nigrisigna, 690
biotic factor, 2043
environmental factor, 2043
flight mill, 690, 2043
Sitodiplosis mosellana, 2043
flight initiation,
DAMs, 2259
insecticide resistance, 2259
walking, 2259
flight mill, 690, 2043
actograph, 690
Autographa nigrisigna, 690
biotic factor, 2043
environmental factor, 2043
flight ability, 690, 2043
Sitodiplosis mosellana, 2043
flight tunnel,
insect behavior, 2144
λ-cyhalothrin, 2144
lethal effect, 2144
sex pheromone, 2144
flonicamid,
cotton aphid, 1430
neonicotinoid, 1430
resistance, 1430
sulfoxaflor, 1430
floral compounds,
dose–response, 1653
electroantennogram, 1653
linalool, 1653
floral traits,
Allium cepa L., 1386
male sterility, 1386
nectar composition, 1386
seed yield, 1386
flowering plant,
conservation biological control, 2327
habitat management, 2327
natural enemy, 2327
Sphaerophoria spp., 2327
fluctuating thermal regime,
alfalfa leafcutting bee, 1081
insect cold storage, 1081
Megachile rotundata, 1081
fluon,
Agrilus planipennis, 2415
green, 2415
multifunnel trap, 2415
purple, 2415
fluorescein sodium salt,
cyantraniliprole, 597
Cyazypyr, 597
imidacloprid, 597
spirotetramat, 597
follicle,
Apolygus lucorum, 1200
diet, 1200
Helicoverpa armigera egg, 1200
ovarian development, 1200
food availability,
Apis mellifera, 2280
honey bee, 2280
hygienic behavior, 2280
seasonal effect, 2280
food-processing facility,
heat treatment, 2247
model, 2247
mortality, 2247
Tribolium castaneum, 2247
foraging behavior,
fipronil, 2160
Reticulitermes, 2160
termite, 2160
termiticide, 2160
forecasting activity,
ambrosia beetle, 1780
bottle trap, 1780
phenological indicator, 1780
trap tree, 1780
forensic entomology,
carrion, 1927
myiasis, 1927
secondary screwworm, 1927
spray-dried blood, 1927
forest pest,
Endoclita signifier, 866
Eucalyptus, 866
life cycle, 866
morphology, 866
Formica sp.,
blueberry, 716
insecticide toxicity, 716
mound-building ants, 716
Fragaria ananassa,
action threshold, 1757
degree-day, 1757
mite-day, 1757
six-spotted thrip, 1757
Frankliniella fusca,
peanut, 587
resistant genotypes, 587
Tospovirus, 587
Fraxinus uhdei,
Agrilus planipennis, 2101
development, 2101
invasive, 2101
laboratory rearing, 2101
freezing, 653, 2433
bed bug, 2433
coffee berry borer, 653
cold tolerance, 2433
green coffee, 653
Hawaii, 653
Hypothenemus hampei, 653
nonchemical control, 2433
supercooling point, 2433
fruit,
cranberry, 2424
ripeness, 2424
spotted wing drosophila, 2424
susceptibility, 2424
fruit firmness,
blueberry, 1767
cultivar, 1767
Drosophila suzukii, 1767
oviposition, 1767
fruit safety,
fruit shelf life, 299
Mangifera indica, 299
Oecophylla smaragdina, 299
soluble sugar content, 299
fruit set,
apical-meristem injury, 1339
cranberry, 1339
side-shoot, 1339
tipworm, 1339
fruit shelf life,
fruit safety, 299
Mangifera indica, 299
Oecophylla smaragdina, 299
soluble sugar content, 299
full-count,
mint, 1707
presence–absence, 1707
twospotted spider mite, 1707
fumigation, 499, 2267, 2341
lemon, 2341
monitoring, 499
nitric oxide, 2267
orange, 2341
pheromone trap, 499
phytosanitary, 2267, 2341
postharvest pest control, 2267
quarantine, 2267
red flour beetle, 499
sulfuryl fluoride, 499
Vapormate, 2341
fumonisin,
borer damage, 2241
Fusarium verticillioides, 2241
Sesamia nonagrioides, 2241
Zea mays, 2241

Fusarium,
integrated pest management (IPM), 1218
monitoring, 1218
trapping, 1218
vector, 1218
Fusarium verticillioides,
borer damage, 2241
fumonisin, 2241
Sesamia nonagrioides, 2241
Zea mays, 2241

G

Galendromus occidentalis,
biological control, 1996
inundative release, 1996
Phytoseiidae, 1996
Tetranychidae, 1996
gall,
aphid, 1628
clone dynamics, 1628
development, 1628
gall wasp,
alternating generation, 1747
bullet gall, 1747
Quercus virginiana, 1747
gamma radiation,
Bacillus thuringiensis, 36
kilogray, 36
SDS-PAGE, 36
toxicity, 36
GC-MS,
botanical insecticides, 159
essential oil, 159
scarab larvae, 159
white grub, 159
gel bait,
German cockroach, 945
indoxacarb, 945
insecticide resistance, 945
resistance monitoring, 945
gene expression, 393, 1446
Bemisia tabaci, 1446
CYP4, 393
cytochrome P450 monooxygenase, 393
Diaphorina citri, 393
endogenous reference gene, 1446
protein expression, 393
quantitative PCR, 1446
RNA-seq, 1446
gene flow,
AFLP, 2225
cotton pest, 2225
dispersal, 2225
population structure, 2225
generalist predator, 776, 1126, 2310
biological control, 776, 1126
fitness, 2310
green lacewing, 1126
inundative release, 2310
invasive species, 2310
lethal, 776
mass-rearing, 2310
OP-replacement insecticides, 1126
pear psylla, 776
reduced-risk insecticide, 776, 1126
sublethal effect, 776
generic dose,
irradiation, 2020
Mediterranean fruit fly, 2020
modified atmosphere, 2020
quarantine, 2020
genetic differentiation,
Homoptera, 1004
ISSR, 1004
molecular marker, 1004
Psyllidae, 1004
genetic distance,
COI, 408
phylogenetic analysis, 408
thrip, 408
genetic engineering,
Bt plant, 2151
insecticide reduction, 2151
Lepidoptera, 2151
maize, 2151
genetic sexing,
Anastrepha ludens, 735
sexual behavior, 735
sterile insect technique, 735
Tapachula-7, 735
genetic sexing strain,
Bactrocera cucurbitae, 995
Bactrocera dorsalis, 995
polytene chromosome, 995
Tephritidae, 995
geographical origin,
bee strain, 1958
botanical origin, 1958
10-hydroxy-*trans*-2-decenoic acid, 1958
royal jelly, 1958
geographic information systems,
Diabrotica barberi, 168
Diabrotica virgifera virgifera, 168
economic injury level, 168
integrated pest management (IPM), 168
German cockroach,
gel bait, 945
indoxacarb, 945
insecticide resistance, 945
resistance monitoring, 945
germplasm,
English grain aphid, 1473
resistance, 1473
tolerance, 1473
wheat, 1473
Glenea cantor,
adult starvation, 215
daily reproductive rhythm, 215
lifetime reproductive pattern, 215
sexual maturation, 215
glucosinolate,
Crambe spp., 2557
flea beetle, 2557
Phyllotreta spp., 2557
Sinapis alba, 2557
Glycine max,
defoliation, 830
insect resistance, 830
value added traits, 830
Gossypium hirsutum, 1045, 2077
cotton boll injury, 2077
cotton fleahopper, 1045
insect–plant interaction, 2077
plant bug, 2077
plant resistance, 1045
Pseudatomoscelis seriatus, 1045
Grandevo,
aphid, 257
biorational, 257
Chromobacterium subtsugae, 257
Curculio caryae, 257
granule formulation,
blacklegged tick, 57
Ixodes scapularis, 57
Metarhizium brunneum, 57
microsclerotia, 57
grape, 482, 905, 1716
Acerophagus, 1716
economic injury level, 905
Ferrisia gilli, 1716
integrated pest management (IPM), 905
mealybug, 482, 1716
monitoring, 482
Paralobesia viteana, 905
pheromone, 482
Vitis, 1716
Washington state vineyard, 482
grape pest management,
Abagrotis, 2135
cutworm, 2135
insecticide toxicity, 2135
grapevine yellows,
Auchenorrhyncha, 604
feeding-medium assay, 604
phytoplasma-insect relationships, 604
phytoplasma seasonal variation, 604
grasshopper bait,
Alaska, 669
Conservation Reserve Program, 669

high-latitude agriculture, 669
Insecta, 669
green,
Agrilus planipennis, 2415
fluon, 2415
multifunnel trap, 2415
purple, 2415
green coffee,
coffee berry borer, 653
freezing, 653
Hawaii, 653
Hypothenemus hampei, 653
greenhouse biocontrol,
Diglyphus isaea, 1590
greenhouse pest management, 1590
leafminer biocontrol, 1590
safe pesticide, 1590
greenhouse pest management,
Diglyphus isaea, 1590
greenhouse biocontrol, 1590
leafminer biocontrol, 1590
safe pesticide, 1590
green lacewing,
biological control, 1126
generalist predator, 1126
OP-replacement insecticides, 1126
reduced-risk insecticide, 1126
grinding,
debarking, 235
firewood, 235
oak mortality, 235
solarization, 235
grooming behavior,
mite fall, 566
Russian honey bee, 566
Varroa destructor, 566
Varroa resistance, 566
group size,
Coptotermes formosanus, 2630
fipronil, 2630
horizontal transfer, 2630
Gryon japonicum,
egg parasitism, 1973
kairomone, 1973
Ooencyrtus nezarae, 1973
soybean, 1973

H

habitat fidelity,
one-choice, 1932
spotted wing drosophila, 1932
4-way olfactometer, 1932
habitat management,
conservation biological control, 2327
flowering plant, 2327
natural enemy, 2327
Sphaerophoria spp., 2327
Halyomorpha halys,
native stink bugs, 1317
soybean, 1317
thresholds, 1317
harborage preference,
Blattella asahinai, 322
mulch, 322
toxicity, 322
harlequin bug,
choice test, 283
cole crop, 283
Murgantia histrionica, 283
trap crop, 283
harmonization plan,
insecticide, 1190
neonicotinoid, 1190
Popillia japonica, 1190
scarab, 1190
Hawaii,
coffee berry borer, 653
freezing, 653
green coffee, 653
Hypothenemus hampei, 653
hay drying,
Mayetiola destructor (Say), 1164
puparial survival, 1164
quarantine control, 1164
systems approach, 1164
heat treatment,
food-processing facility, 2247
model, 2247
mortality, 2247
Tribolium castaneum, 2247
Helicoverpa armigera, 375, 450, 614, 708, 2514
Bt cotton, 375
Bt toxin, 614
circadian clock, 450
clock genes, 450
Cry1Ac resistance, 2514
cryptochrome, 450
emamectin benzoate, 375
fenvalerate, 375
lifespan, 614
mating history, 708
nonbinding site, 2514
photoreceptor, 450
phoxim, 375
polyandry, 708
proteinases, 2514
reproductive performance, 614
sperm competition, 708
sperm transfer, 708
Spodoptera exigua, 614
Helicoverpa armigera egg,
Apolygus lucorum, 1200
diet, 1200
follicle, 1200
ovarian development, 1200
Helicoverpa zea,
Bacillus thuringiensis, 1250
pheromone trap, 1250
transgenic corn, 1250
Hemiptera vector,
dispersal, 2296
plant community, 2296
vineyard agroecosystem, 2296
hemlock woolly adelgid,
canopy translocation, 2399
neonicotinoid, 2399
systemic insecticide, 2399
Tsuga canadensis, 2399
hemolymph protein,
artificial diet, 2364
Attacus atlas, 2364
Heterorhabditis, 622, 1112
Diabrotica barberi, 622
Diabrotica virgifera virgifera, 622
entomopathogenic nematode, 1112
Metarhizium, 622
peanut, 1112
Steinernema, 622, 1112
white grub, 1112
Heterorhabditis bacteriophora,
Beauveria bassiana, 2072
burrower bug, 2072
chlorpyrifos, 2072
high-latitude agriculture,
Alaska, 669
Conservation Reserve Program, 669
grasshopper bait, 669
Insecta, 669
Histeridae,
Cerambycidae, 206
Cleridae, 206
Scolytinae, 206
Trogossitidae, 206
Homoptera,
genetic differentiation, 1004
ISSR, 1004
molecular marker, 1004
Psyllidae, 1004
honey bee, 558, 1535, 1553, 2280
amitraz, 558
Apis mellifera, 2280
fat body, 1553
food availability, 2280
hygienic behavior, 2280
miticides, 558
nutritional status, 1553
pollen diets, 1553
protein, 1553
screening protocols, 1535
seasonal effect, 2280
surveillance, 1535
tau-fluvalinate, 558
Tropilaelaps mite, 1535
Varroa destructor, 558

horizontal transfer,
Coptotermes formosanus, 2630
fipronil, 2630
group size, 2630
horticulture,
antagonism, 247
bioassay, 247
pest management, 247
synergism, 247
host age,
biological control, 633
egg-parasitoid, 633
western conifer seed bug, 633
host-feeding,
biological control, 1638
integrated pest management (IPM), 1638
life table, 1638
parasitoid behavior, 1638
host plant, 1404, 1414
Bemisia tabaci, 1414
desert environment, 1404
IGRs, 1414
insecticide resistance, 1404
neonicotinoid, 1414
resistance, 1414
systemic uptake bioassay, 1404
host-plant resistance, 1036, 1479, 1871, 2577
adult damage, 1871
Aphis glycines, 1036, 1479
biotype, 1479
Brachiaria, 1871
nutritional index, 2577
preference, 2577
Rag1 gene, 2577
soybean, 1479, 2577
spittlebug, 1871
tolerance, 1036
host plant suitability,
Aphididae, 683
electrical penetration graph, 683
Miscanthus sacchariflorus, 683
phytovirus, 683
host recognition,
behavior, 2354
Diabrotica virgifera virgifera, 2354
larva, 2354
Zea mays, 2354
host susceptibility,
Bactrocera tryoni, 883
citrus, 883
field infestation, 883
host switch,
cereal crops, 1294
cultural control, 1294
English grain aphid, 1294
life table, 1294
host use,
boll weevil, 437
COI DNA, 437
cotton pest, 437
DNA barcode, 437
huanglongbing,
Asian citrus psyllid, 25
Candidatus Liberibacter asiaticus, 25
citrus greening, 25
Diaphorina citri, 25
10-hydroxy-*trans*-2-decenoic acid,
bee strain, 1958
botanical origin, 1958
geographical origin, 1958
royal jelly, 1958
hygienic behavior,
Apis mellifera, 2280
food availability, 2280
honey bee, 2280
seasonal effect, 2280
Hypera postica,
alfalfa weevil, 426
phylogenetic, 426
Hypothenemus hampei,
coffee berry borer, 653
freezing, 653
green coffee, 653
Hawaii, 653

I

identification,
antifungal activity, 43
Bacillus subtilis, 43
Blattella germanica, 43
intestinal flora, 43
IGRs,
Bemisia tabaci, 1414
host plant, 1414
neonicotinoid, 1414
resistance, 1414
imidacloprid, 400, 597, 807, 874, 1064
acephate, 807
Acer platanoides, 874
aldicarb, 807
binary combinations, 1064
cyantraniliprole, 597
Cyazypyr, 597
cyfluthrin, 400, 1064
distribution, 874
fluorescein sodium salt, 597
metamidophos, 400
Myzus persicae, 400
neonicotinoid, 874
pirimicarb, 400
soil-applied insecticides, 807
spirotetramat, 597
surface treatment, 1064
thiamethoxam, 807
translocation, 874
immunohistochemistry,
Aedes albopictus, 1098
Bacillus thuringiensis, 1098
ELISA, 1098
receptor binding, 1098
incidence,
corn leafhopper, 1574
mollicutes, 1574
overwintering, 1574
pest, 1574
indoxacarb, 945, 1423
baseline toxicity, 1423
gel bait, 945
German cockroach, 945
insecticide resistance, 945
metaflumizone, 1423
Plutella xylostella, 1423
resistance monitoring, 945
infestation,
bed bug, 347
Cimex lectularius, 347
control, 347
dust, 347
injury,
blackberry, 912
Euschistus spp., 912
monitoring, 912
stink bug, 912
injury equivalency,
feeding guild, 837
Oryza sativa, 837
rice, 837
rice planthopper, 837
insect, 382, 1179, 1659
biosecurity, 1179
detection capacity, 1659
environmental risk assessment, 1659
mite, 1179
phytosanitary, 1179
plant quarantine, 1179
population genetic, 382
statistical power, 1659
stochastic processes, 382
transgenic crop, 1659
Insecta,
Alaska, 669
Conservation Reserve Program, 669
grasshopper bait, 669
high-latitude agriculture, 669
insect behavior, 1463, 2144
antibiosis, 1463
antixenosis, 1463
crane fly, 1463
endophytes, 1463
flight tunnel, 2144
λ-cyhalothrin, 2144
lethal effect, 2144
sex pheromone, 2144

insect cold storage,
alfalfa leafcutting bee, 1081
fluctuating thermal regime, 1081
Megachile rotundata, 1081
insect colony effect,
production, 50
SeMNPV, 50
Spexit, 50
Vir-ex, 50
insect damage,
cotton, 1209
lint yield, 1209
Lygus hesperus, 1209
plant compensation, 1209
insect–disease complex,
chemical control, 2286
exotic insect, 2286
Lauraceae, 2286
redbay ambrosia beetle, 2286
insecticide, 747, 1190, 1863
Coptotermes formosanus, 1863
differentially expressed gene, 1863
egg, 747
embryology, 747
fall armyworm, 747
harmonization plan, 1190
morphology, 747
neonicotinoid, 1190
Popillia japonica, 1190
scarab, 1190
insecticide efficacy,
invasive species, 2448
occasional invader, 2448
insecticide formulation,
insecticide susceptibility, 2530
insecticide toxicity, 2530
resistance, 2530
insecticide reduction,
Bt plant, 2151
genetic engineering, 2151
Lepidoptera, 2151
maize, 2151
insecticide resistance, 366, 939, 945, 954, 959, 1404, 1832, 1855, 2167, 2208, 2259, 2491
Aedes aegypti, 959
beet armyworm, 1855
Brevicoryne brassicae, 954
Chilo suppressalis, 1832
codling moth, 939
Colorado potato beetle, 2491
cross-resistance, 959, 2167
DAMs, 2259
desert environment, 1404
diagnostic concentration, 939
diapause, 2491
dieldrin, 2167
dynamics, 1832
economic, 366
fipronil, 2167
fitness costs, 366
flight initiation, 2259
gel bait, 945
German cockroach, 945
host plant, 1404
indoxacarb, 945
integrated pest management (IPM), 2491
lesser grain borer, 2208
malaria, 366
neonate, 939
neonicotinoid, 954
organophosphate, 954
phenology, 2491
phosphine resistance management, 2208
pyrethroid, 954, 959
Rdl gene, 2167
red flour beetle, 2208
regression analysis, 1832
resistance monitoring, 945, 1855
stored product, 2208
systemic uptake bioassay, 1404
vector control, 366
walking, 2259
insecticide susceptibility,
insecticide formulation, 2530
insecticide toxicity, 2530
resistance, 2530
insecticide timing,
broccoli, 267
cabbage, 267
integrated pest management (IPM), 267
invasive species, 267
insecticide toxicity, 353, 716, 2135, 2530
Abagrotis, 2135
Bactrocera dorsalis (Hendel), 353
blueberry, 716
cutworm, 2135
environmental factor, 353
Formica sp., 716
grape pest management, 2135
insecticide formulation, 2530
insecticide susceptibility, 2530
mound-building ants, 716
resistance, 2530
insect infestation,
mineral, 1058
Sitophilus granarius, 1058
storage, 1058
vitamin, 1058
insect pest,
booklouse, 491
Liposcelis, 491
Psocoptera, 491
stored product, 491
insect–plant interaction, 1134, 1240, 2077
biological control, 1134
cabbage, 1134
cotton boll injury, 2077
Gossypium hirsutum, 2077
kale, 1134
N availability, 1240
parasitoid community, 1134
plant bug, 2077
Spodoptera exigua, 1240
Zea mays, 1240
insect pollinator,
Echinacea angustifolia, 2055
pollen germination, 2055
pollen-tube quantification, 2055
shrivelled (retracted) styles, 2055
insect resistance,
defoliation, 830
Glycine max, 830
value added traits, 830
insect resistance evaluation,
Cry1Ab/Cry2Aj protein, 2548
Cry1Ab/Vip3DA protein, 2548
Ostrinia furnacalis, 2548
transgenic maize, 2548
insect resistance management, 1525, 1819, 2473, 2506
Bacillus thuringiensis, 1525, 2506
Bt cotton, 1819
Bt resistance, 2506
Callosobruchus maculatus, 2473
cowpea, 2473
Diabrotica virgifera virgifera, 2506
field-evolved resistance, 1525
refuge, 1525
simulation, 2473
tobacco budworm, 1819
transgenic crop, 2473
instar determination method,
kernel density estimation, 800
nitrogen, 800
Spodoptera exigua, 800
integrated pest management (IPM), 168, 267, 648, 905, 1218, 1302, 1638, 1739, 1920, 2126, 2491
Aphis glycines, 1302
astigmata, 1920
biological control, 1638
broccoli, 267
cabbage, 267
Colorado potato beetle, 2491
Diabrotica barberi, 168
Diabrotica virgifera virgifera, 168
diapause, 2491
disinfestation, 1920

dispersal, 1920
economic injury level, 168, 905
entomopathogenic nematode, 648
ethyl acetate, 1739
Fusarium, 1218
geographic information systems, 168
grape, 905
host-feeding, 1638
insecticide resistance, 2491
insecticide timing, 267
invasive species, 267
kairomone, 1739
life table, 1638
monitoring, 1218
natural insecticide, 1920
Paralobesia viteana, 905
parasitoid behavior, 1638
phenology, 2491
red palm weevil, 1739
selective insecticide, 2126
Steinernema carpocapsae, 648
Steinernematidae, 648
systemics, 2126
Tenebrionidae, 648
thiamethoxam, 1302
trap, 1739
trapping, 1218
vector, 1218
water volume, 2126
internal transcribed spacer,
cryptic species, 463
cytochrome *c* oxidase subunit I, 463
DNA-based identification, 463
Tetranychus, 463
intestinal flora,
antifungal activity, 43
Bacillus subtilis, 43
Blattella germanica, 43
identification, 43
intrinsic rate of increase, 675, 2177, 2595
actuarial mathematics, 675
alternative host plant, 2595
Anarsia lineatella, 675
bio-insecticide, 2177
biotype, 2595
demographic parameter, 675
Diuraphis noxia, 2595
finite rate of increase, 2177
life table, 675
net reproductive rate, 2177
population parameter, 2177
Russian wheat aphid, 2595
inundative release, 1996, 2310
biological control, 1996
fitness, 2310
Galendromus occidentalis, 1996
generalist predator, 2310
invasive species, 2310
mass-rearing, 2310
Phytoseiidae, 1996
Tetranychidae, 1996
invasive, 1145, 2101
Agrilus planipennis, 2101
barcode, 2275
cockroach, 2275
development, 2101
establishment, 1145
Fraxinus uhdei, 2101
laboratory rearing, 2101
natural enemy introduction, 1145
parasitoid release, 1145
Periplaneta japonica, 2275
pest, 2275
wood borers, 1145
invasive ants,
ant, 1545
control, 1545
eradication, 1545
treatment, 1545
invasive carpophagous species,
mortality, 2349
wine grape, 2349
winery waste, 2349
invasive forest pest,
beech bark disease, 1324
Cryptococcus fagisuga, 1324
Fagus grandifolia, 1324
invasive species, 150, 267, 311, 1355, 1676, 2216, 2310, 2428, 2448
Bemisia tabaci, 1355
Blatta orientalis, 2428
blattidae, 2428
breeding system, 2216
broccoli, 267
brown marmorated stink bug, 150
cabbage, 267
fitness, 2310
generalist predator, 2310
insecticide efficacy, 2448
insecticide timing, 267
inundative release, 2310
integrated pest management (IPM), 267
mass-rearing, 2310
Megacopta cribraria, 1676
movement, 150
neoteny, 2216
occasional invader, 2448
oriental cockroach, 2428
pesticide, 150
population genetic, 1355
soybean, 1676
species distribution model, 311
sublethal, 150
subterranean termite, 311
termite, 2216
urban pest, 311
invasive species ecology,
degree-day, 1495
distribution, 1495
survey, 1495
invertebrate,
agro-ecosystems, 756
mortality, 756
nontarget, 756
pesticide, 756
ionizing radiation,
phytosanitary treatment, 525
phytosanitation, 525
quarantine pest, 525
Ipomoea,
Cylas formicarius, 1456
Diabrotica, 1456
plant resistance, 1456
white grub, 1456
ipsdienol,
α-pinene, 1684
ethanol, 1684
ipsenol, 1684
kairomone, 1684
ipsenol,
α-pinene, 1684
ethanol, 1684
ipsdienol, 1684
kairomone, 1684
irradiation, 641, 2020, 2035
generic dose, 2020
medfly, 641
Mediterranean fruit fly, 2020
modified atmosphere, 2020
phytosanitary treatment, 2035
postharvest, 2035
probiotic, 641
quarantine, 2020, 2035
sperm, 641
sterile insect technique, 641
systems approach, 2035
Isoptera,
Coptotermes formosanus, 2442
mark–recapture protocol, 2442
probability density function, 2442
Rhinotermitidae, 2442
ISSR,
genetic differentiation, 1004
Homoptera, 1004
molecular marker, 1004
Psyllidae, 1004
ITS,
PCR-restriction fragment-length polymorphism, 661
ribosomal DNA, 661
species identification, 661
Tetranychus, 661
Ixodes scapularis,
blacklegged tick, 57
granule formulation, 57
Metarhizium brunneum, 57
microsclerotia, 57

J

jackknife,
bootstrap, 1
Chrysopa pallens, 1
life table, 1
population projection, 1
Japanese beetle,
corn, 2048
Popillia japonica, 2048
silk clipping, 2048
Juniperus oxycedrus ssp. *oxycedrus*,
essential oil, 1902
silica gel, 1902
Sitophilus oryzae, 1902
stored wheat, 1902
Tribolium confusum, 1902

K

kairomone, 1684, 1739, 1973
α-pinene, 1684
egg parasitism, 1973
ethanol, 1684
ethyl acetate, 1739
Gryon japonicum, 1973
integrated pest management (IPM), 1739
ipsdienol, 1684
ipsenol, 1684
Ooencyrtus nezarae, 1973
red palm weevil, 1739
soybean, 1973
trap, 1739
kale,
biological control, 1134
cabbage, 1134
insect–plant interaction, 1134
parasitoid community, 1134
kernel density estimation,
instar determination method, 800
nitrogen, 800
Spodoptera exigua, 800
kilogray,
Bacillus thuringiensis, 36
gamma radiation, 36
SDS-PAGE, 36
toxicity, 36
knockdown resistance,
Culex pipiens pallens, 360
pyrethroid insecticide resistance, 360
susceptibility, 360

L

laboratory rearing,
Agrilus planipennis, 2101
development, 2101
Fraxinus uhdei, 2101
invasive, 2101
λ-cyhalothrin,
flight tunnel, 2144
insect behavior, 2144
lethal effect, 2144
sex pheromone, 2144
LAMP,
Bactericera cockerelli, 1440
Candidatus Liberibacter solanacearum, 1440
pathogen detection, 1440
Zebra chip, 1440
larva,
behavior, 2354
Diabrotica virgifera virgifera, 2354
host recognition, 2354
Zea mays, 2354
larval survival,
economic injury level, 1274
economic threshold, 1274
Striacosta albicosta, 1274
western bean cutworm, 1274
Lauraceae,
chemical control, 2286
exotic insect, 2286
insect–disease complex, 2286
redbay ambrosia beetle, 2286
laurel wilt,
Persea borbonia, 1693
Raffaelea lauricola, 1693
redbay ambrosia beetle, 1693
leaf-cutting ant,
Atta sexdens rubropilosa, 742
castor oil, 742
physic nut, 742
seed lipid, 742
leaf disk transmission assay,
semipersistent, 16
vector, 16
leafminer biocontrol,
Diglyphus isaea, 1590
greenhouse biocontrol, 1590
greenhouse pest management, 1590
safe pesticide, 1590
lemon,
fumigation, 2341
orange, 2341
phytosanitary, 2341
Vapormate, 2341
Lepidoptera,
Bt plant, 2151
genetic engineering, 2151
insecticide reduction, 2151
maize, 2151
Leptoglossus occidentalis,
cone abortion, 229
Ernobius spp., 229
nut production, 229
stone pine, 229
lesser grain borer,
insecticide resistance, 2208
phosphine resistance management, 2208
red flour beetle, 2208
stored product, 2208
lethal,
biological control, 776
generalist predator, 776
pear psylla, 776
reduced-risk insecticide, 776
sublethal effect, 776
lethal effect,
flight tunnel, 2144
insect behavior, 2144
λ-cyhalothrin, 2144
sex pheromone, 2144
lettuce crop,
aphid damage, 891
economic threshold, 891
edibility, 891
levels of infestation,
Bactrocera latifrons, 1648
Capsicum annuum, 1648
cultivar, 1648
Malaysian fruit fly, 1648
Liberibacter,
acquisition access period, 1964
potato, 1964
potato psyllid, 1964
tomato, 1964
life cycle,
Endoclita signifier, 866
Eucalyptus, 866
forest pest, 866
morphology, 866
life history,
acquisition, 1878
plant virus, 1878
resistance, 1878
transmission, 1878
lifespan,
Bt toxin, 614
Helicoverpa armigera, 614
reproductive performance, 614
Spodoptera exigua, 614
life table, 1, 533, 675, 1294, 1638, 2118, 2371
actuarial mathematics, 675
Anarsia lineatella, 675
apple, 2118
biological control, 1638, 2371
bootstrap, 1
cereal crops, 1294
Chrysopa pallens, 1
cultural control, 1294
demographic parameter, 675
development, 2118
English grain aphid, 1294
fecundity, 2371
first-generation, 533
host-feeding, 1638

host switch, 1294
integrated pest management (IPM), 1638
intrinsic rate of increase, 675
jackknife, 1
Megacopta cribraria, 533
nontarget effect, 2371
Panonychus ulmi, 2118
parasitoid behavior, 1638
population projection, 1
rearing, 533
reproduction, 2118
soybean, 533
twospotted spider mite, 2371
lifetime reproductive pattern,
adult starvation, 215
daily reproductive rhythm, 215
Glenea cantor, 215
sexual maturation, 215
light brown apple moth,
phosphine, 1613
phytosanitary, 1613
postharvest pest control, 1613
quarantine treatment, 1613
light reflectance,
antixenosis, 2602
Brassica oleracea, 2602
resistance, 2602
Thrips tabaci, 2602
linalool,
dose–response, 1653
electroantennogram, 1653
floral compounds, 1653
lint yield,
cotton, 1209
insect damage, 1209
Lygus hesperus, 1209
plant compensation, 1209
Liposcelis, 419, 491
booklouse, 491
insect pest, 491
molecular identification, 419
morphological characteristics, 419
phylogenetic relationships, 419
Psocoptera, 491
SEM, 419
stored product, 491
Liposcelis bostrychophila,
Epimedium pubescens, 513
(R)-carvone, 513
repellency, 513
Tribolium castaneum, 513
Listronotus maculicollis,
Beauveria bassiana, 2322
Curculionidae, 2322
neonicotinoid, 2322
litchi,
litchi erineum mite, 2183
predatory mite, 2183
selectivity, 2183
litchi erineum mite,
litchi, 2183
predatory mite, 2183
selectivity, 2183
Loberus testaceus,
Erotylidae, 305
Europs, 305
Nitidulidae, 305
Omatomidae, 305
lodgepole pine,
antiaggregant, 221
mountain pine beetle, 221
semiochemical, 221
verbenone, 221
lower developmental threshold,
bootstrap percentile, 107
Performance-2 model, 107
sum of effective temperatures, 107
upper developmental threshold, 107
low oxygen,
apple maggot, 1173
controlled atmosphere, 1173
Diptera: Tephritidae, 1173
low pressure, 1173
low pressure,
apple maggot, 1173
controlled atmosphere, 1173
Diptera: Tephritidae, 1173
low oxygen, 1173
lufenuron,
carcass-burying behavior, 1812
Coptotermes formosanus, 1812
disease resistance, 1812
Pseudomonas aeruginosa, 1812
Lygus,
cotton, 1286
ecoinformatics, 1286
farmer decision-making, 1286
observational studies, 1286
Lygus hesperus,
cotton, 1209
insect damage, 1209
lint yield, 1209
plant compensation, 1209
Lygus lineolaris,
cotton, 2378
cultural control, 2378

M

maize, 1234, 1669, 2151
Bt plant, 2151
dead-heart plants, 1669
Diabrotica undecimpunctata howardi, 1669
European corn borer, 1234
genetic engineering, 2151
insecticide reduction, 2151
Lepidoptera, 2151
Mediterranean corn borer, 1234
plant development, 1234
resistance, 1234
seedling damage, 1669
southern corn rootworm, 1669
malaria,
economic, 366
fitness costs, 366
insecticide resistance, 366
vector control, 366
Malaysian fruit fly,
Bactrocera latifrons, 1648
Capsicum annuum, 1648
cultivar, 1648
levels of infestation, 1648
male sterility,
Allium cepa L., 1386
floral traits, 1386
nectar composition, 1386
seed yield, 1386
Malus spectabilis,
apple maggot, 1511
microsatellites, 1511
Rhagoletis indifferens, 1511
western cherry fruit fly, 1511
Mangifera indica,
fruit safety, 299
fruit shelf life, 299
Oecophylla smaragdina, 299
soluble sugar content, 299
mark–recapture protocol,
Coptotermes formosanus, 2442
Isoptera, 2442
probability density function, 2442
Rhinotermitidae, 2442
mass-rearing,
fitness, 2310
generalist predator, 2310
inundative release, 2310
invasive species, 2310
mating assessment,
Mediterranean fruit fly, 2541
PCR-restriction fragment-length polymorphism, 2541
stored sperm identification, 2541
mating behavioral sequence,
courtship, 1373
mating success, 1373
sexual selection, 1373
mating history,
Helicoverpa armigera, 708
polyandry, 708
sperm competition, 708
sperm transfer, 708
mating success,
courtship, 1373
mating behavioral sequence, 1373
sexual selection, 1373

Matsucoccidae,
eastern white pine, 2391
Matsucoccus macrocicatrices, 2391
Pinus strobus, 2391
southeastern United States, 2391
Matsucoccus macrocicatrices,
eastern white pine, 2391
Matsucoccidae, 2391
Pinus strobus, 2391
southeastern United States, 2391
maximum temperature,
brown marmorated stink bug, 2141
overwintering adult, 2141
population density, 2141
Mayetiola destructor (Say),
hay drying, 1164
Mayetiola destructor (Say), 1164
puparial survival, 1164
quarantine control, 1164
systems approach, 1164
mealybug, 482, 1716
Acerophagus, 1716
Ferrisia gilli, 1716
grape, 482, 1716
monitoring, 482
pheromone, 482
Vitis, 1716
Washington state vineyard, 482
medfly,
irradiation, 641
probiotic, 641
sperm, 641
sterile insect technique, 641
median lethal dose,
biological control, 1582
median lethal time, 1582
minimum inhibitory concentration, 1582
total hemocyte count, 1582
median lethal time,
biological control, 1582
median lethal dose, 1582
minimum inhibitory concentration, 1582
total hemocyte count, 1582
Medicago sativa,
alfalfa-seed production, 576
Megachile rotundata, 576
Megachilidae, 576
pollination, 576
Mediterranean corn borer,
European corn borer, 1234
maize, 1234
plant development, 1234
resistance, 1234
Mediterranean fruit fly, 1772, 2020, 2336, 2541
azadirachtin, 1772
cold treatment, 2336
generic dose, 2020
irradiation, 2020
mating assessment, 2541
modified atmosphere, 2020
neem, 1772
PCR-restriction fragment-length polymorphism, 2541
peach fruit fly, 2336
phytosanitary treatment, 2336
quarantine, 2020
quarantine treatment, 2336
South American fruit fly, 1772
stored sperm identification, 2541
toxic bait, 1772
Megachile rotundata, 543, 576, 1081, 1089
alfalfa, 543
alfalfa leafcutting bee, 1081
alfalfa-seed production, 576
diapause, 1089
fluctuating thermal regime, 1081
insect cold storage, 1081
Medicago sativa, 576
Megachilidae, 543, 576
metabolic rate, 1089
pollination, 543, 576
reproduction, 543
respiration, 1089
thermal stress, 1089
Megachilidae, 543, 576
alfalfa, 543
alfalfa-seed production, 576
Medicago sativa, 576
Megachile rotundata, 543, 576
pollination, 543, 576
reproduction, 543
Megacopta cribraria, 533, 1676
first-generation, 533
invasive species, 1676
life table, 533
rearing, 533
soybean, 533, 1676
metabolic rate,
diapause, 1089
Megachile rotundata, 1089
respiration, 1089
thermal stress, 1089
metaflumizone,
baseline toxicity, 1423
indoxacarb, 1423
Plutella xylostella, 1423
metamidophos,
cyfluthrin, 400
imidacloprid, 400
Myzus persicae, 400
pirimicarb, 400
Metarhizium,
Diabrotica barberi, 622
Diabrotica virgifera virgifera, 622
Heterorhabditis, 622
Steinernema, 622
Metarhizium anisopliae strain F52,
acquisition, 277
Bombus terrestris, 277
miniature dispenser, 277
worker mortality, 277
Metarhizium brunneum,
blacklegged tick, 57
granule formulation, 57
Ixodes scapularis, 57
microsclerotia, 57
methoprene,
economic analysis, 1503
esfenvalerate, 1503
Plodia interpunctella, 1503
methyl bromide,
Epiphyas postvittana, 1155
postharvest fumigation, 1155
sorption kinetics, 1155
methyl eugenol,
monitoring traps, 727
polymer matrix, 727
spinosad, 727
toxicant, 727
microbial control,
biological control, 1118
crude extract, 1118
olive fruit fly, 1118
secondary metabolite, 1118
microsatellites,
apple maggot, 1511
Malus spectabilis, 1511
Rhagoletis indifferens, 1511
western cherry fruit fly, 1511
microsclerotia,
blacklegged tick, 57
granule formulation, 57
Ixodes scapularis, 57
Metarhizium brunneum, 57
mineral,
insect infestation, 1058
Sitophilus granarius, 1058
storage, 1058
vitamin, 1058
miniature dispenser,
acquisition, 277
Bombus terrestris, 277
Metarhizium anisopliae strain F52, 277
worker mortality, 277
minimum inhibitory concentration,
biological control, 1582
median lethal dose, 1582
median lethal time, 1582
total hemocyte count, 1582

mint,
full-count, 1707
presence–absence, 1707
twospotted spider mite, 1707
Miscanthus sacchariflorus,
Aphididae, 683
electrical penetration graph, 683
host plant suitability, 683
phytovirus, 683
mite,
biosecurity, 1179
insect, 1179
phytosanitary, 1179
plant quarantine, 1179
mite-day,
action threshold, 1757
degree-day, 1757
Fragaria ananassa, 1757
six-spotted thrip, 1757
mite fall,
grooming behavior, 566
Russian honey bee, 566
Varroa destructor, 566
Varroa resistance, 566
miticides,
amitraz, 558
honey bee, 558
tau-fluvalinate, 558
Varroa destructor, 558
model,
food-processing facility, 2247
heat treatment, 2247
mortality, 2247
Tribolium castaneum, 2247
modified atmosphere,
generic dose, 2020
irradiation, 2020
Mediterranean fruit fly, 2020
quarantine, 2020
molecular identification,
Liposcelis, 419
morphological characteristics, 419
phylogenetic relationships, 419
SEM, 419
molecular marker, 1004, 2457
F1538I mutation, 2457
fenpropathrin, 2457
genetic differentiation, 1004
Homoptera, 1004
ISSR, 1004
Psyllidae, 1004
Tetranychus cinnabarinus, 2457
voltage-gate sodium channel, 2457
mollicutes,
corn leafhopper, 1574
incidence, 1574
overwintering, 1574
pest, 1574
monitoring, 482, 499, 844, 912, 1218
blackberry, 912
clover, 844
Coleophora deauratella, 844
Euschistus spp., 912
fumigation, 499
Fusarium, 1218
grape, 482
injury, 912
integrated pest management (IPM), 1218
mealybug, 482
pheromone, 482
pheromone-baited traps, 844
pheromone trap, 499
red flour beetle, 499
stink bug, 912
sulfuryl fluoride, 499
trapping, 1218
vector, 1218
Washington state vineyard, 482
monitoring traps,
methyl eugenol, 727
polymer matrix, 727
spinosad, 727
toxicant, 727
morning glory,
aphid, 1566
potyviruses, 1566
sweet potato, 1566
morphological characteristics,
Liposcelis, 419
molecular identification, 419
phylogenetic relationships, 419
SEM, 419
morphology, 747, 866
egg, 747
embryology, 747
Endoclita signifier, 866
Eucalyptus, 866
fall armyworm, 747
forest pest, 866
insecticide, 747
life cycle, 866
mortality, 756, 786, 1911, 2247, 2349
agro-ecosystems, 756
chlorpyrifos-methyl plus deltamethrin, 1911
damage, 786
field strains, 1911
food-processing facility, 2247
heat treatment, 2247
invasive carpophagous species, 2349
invertebrate, 756
model, 2247
nontarget, 756
pesticide, 756
potato, 786
recovery, 786
spinosad, 1911
stored-grain insects, 1911
Tribolium castaneum, 2247
wine grape, 2349
winery waste, 2349
wireworm, 786
mound-building ants,
blueberry, 716
Formica sp., 716
insecticide toxicity, 716
mountain pine beetle,
antiaggregant, 221
lodgepole pine, 221
semiochemical, 221
verbenone, 221
movement,
brown marmorated stink bug, 150
invasive species, 150
pesticide, 150
sublethal, 150
mulch,
Blattella asahinai, 322
harborage preference, 322
toxicity, 322
multifunnel trap, 2407, 2415
Agrilus planipennis, 2415
detection tools, 2407
fluon, 2415
green, 2415
prism trap, 2407
purple, 2415
Sabic green, 2407
Sabic purple, 2407
multivariate statistical analysis,
principal component analysis, 1053
tritrophic interaction, 1053
VOCs, 1053
Murgantia histrionica,
choice test, 283
cole crop, 283
harlequin bug, 283
trap crop, 283
Musca domestica,
attractant compounds, 979
vinegar, 979
mushroom,
nonlinear modeling, 115
thermal requirement, 115
variety, 115
myiasis,
carrion, 1927
forensic entomology, 1927
secondary screwworm, 1927
spray-dried blood, 1927
Mymaridae,
biological control, 1979
Cleruchoides noackae, 1979
per capita reproduction, 1979
Thaumastocoris peregrinus, 1979

Myzus persicae, 400, 1699, 1838
alphacypermethrin, 1699
Aphidius colemani, 1699
cabbage, 1699
carbamate, 1838
cyfluthrin, 400
imidacloprid, 400
metamidophos, 400
oilseed rape, 1838
pirimicarb, 400
pyrethroid, 1838
repellent netting, 1699
target site resistance, 1838

N

native species,
cross-resistance, 1029
plant resistance, 1029
wheat pests, 1029
native stink bugs,
Halyomorpha halys, 1317
soybean, 1317
thresholds, 1317
natural enemy,
conservation biological control, 2327
flowering plant, 2327
habitat management, 2327
Sphaerophoria spp., 2327
natural enemy introduction,
establishment, 1145
invasive, 1145
parasitoid release, 1145
wood borers, 1145
natural insecticide,
astigmata, 1920
disinfestation, 1920
dispersal, 1920
integrated pest management (IPM), 1920
natural product,
bed bug, 2522
DEET, 2522
essential oil, 2522
repellent, 2522
natural target pruning,
boring larva, 1602
branch protection zone, 1602
olive pyralid moth, 1602
pruning paint, 1602
N availability,
insect-plant interaction, 1240
Spodoptera exigua, 1240
Zea mays, 1240
nectar composition,
Allium cepa L., 1386
floral traits, 1386
male sterility, 1386
seed yield, 1386
neem,
azadirachtin, 1772
Mediterranean fruit fly, 1772
South American fruit fly, 1772
toxic bait, 1772
neonate,
codling moth, 939
diagnostic concentration, 939
insecticide resistance, 939
neonicotinoid, 874, 954, 1190, 1414, 1430, 2322, 2399
Acer platanoides, 874
Beauveria bassiana, 2322
Bemisia tabaci, 1414
Brevicoryne brassicae, 954
canopy translocation, 2399
cotton aphid, 1430
Curculionidae, 2322
distribution, 874
flonicamid, 1430
harmonization plan, 1190
hemlock woolly adelgid, 2399
host plant, 1414
IGRs, 1414
imidacloprid, 874
insecticide, 1190
insecticide resistance, 954
Listronotus maculicollis, 2322
organophosphate, 954
Popillia japonica, 1190
pyrethroid, 954
resistance, 1414, 1430
scarab, 1190
sulfoxaflor, 1430
systemic insecticide, 2399
translocation, 874
Tsuga canadensis, 2399
neoteny,
breeding system, 2216
invasive species, 2216
termite, 2216
net reproductive rate,
bio-insecticide, 2177
finite rate of increase, 2177
intrinsic rate of increase, 2177
population parameter, 2177
Nilaparvata lugens, 10, 1887
Bt rice, 1887
detoxification enzyme, 1887
ecological fitness, 1887
fecundity, 10
population, 1887
sublethal effect, 10
survival, 10
wing formation, 10
Nitidulidae,
Erotylidae, 305
Europs, 305
Loberus testaceus, 305
Omatomidae, 305
nitric oxide,
fumigation, 2267
phytosanitary, 2267
postharvest pest control, 2267
quarantine, 2267
nitrogen,
instar determination method, 800
kernel density estimation, 800
Spodoptera exigua, 800
nitrogen fertilizer,
biochemistry, 196
Cnaphalocrocis medinalis, 196
elemental magnesium, 196
fecundity, 196
nonbinding site,
Cry1Ac resistance, 2514
Helicoverpa armigera, 2514
proteinases, 2514
nonchemical control,
bed bug, 2433
cold tolerance, 2433
freezing, 2433
supercooling point, 2433
nonlinear modeling,
mushroom, 115
thermal requirement, 115
variety, 115
nonreproducing state,
cotton, 86
tropical population, 86
nontarget,
agro-ecosystems, 756
invertebrate, 756
mortality, 756
pesticide, 756
nontarget effect,
biological control, 2371
fecundity, 2371
life table, 2371
twospotted spider mite, 2371
nonvector herbivore,
Bemisia tabaci, 924
fitness, 924
tomato spotted wilt virus, 924
northern corn rootworm,
overwintering, 1619
population variation, 1619
nursery stock,
Cicadellidae, 2027
Dracaena marginata, 2027
offshore preclearance program, 2027
size restriction, 2027
nut production,
cone abortion, 229
Ernobius spp., 229
Leptoglossus occidentalis, 229
stone pine, 229
nutritional index, 1018, 2577
digestive enzymes, 1018
host-plant resistance, 2577
Phthorimaea operculella, 1018
preference, 2577
potato germplasms, 1018
Rag1 gene, 2577
soybean, 2577

nutritional status,
fat body, 1553
honey bee, 1553
pollen diets, 1553
protein, 1553
nylon 12,
Coptotermes, 1395
plastic cable sheathings, 1395
polyethylene, 1395
subterranean termite, 1395

O

oak mortality,
debarking, 235
firewood, 235
grinding, 235
solarization, 235
observational studies,
cotton, 1286
ecoinformatics, 1286
farmer decision-making, 1286
Lygus, 1286
occasional invader,
insecticide efficacy, 2448
invasive species, 2448
Oecophylla smaragdina,
fruit safety, 299
fruit shelf life, 299
Mangifera indica, 299
soluble sugar content, 299
offshore preclearance program,
Cicadellidae, 2027
Dracaena marginata, 2027
nursery stock, 2027
size restriction, 2027
oilseed rape,
carbamate, 1838
Myzus persicae, 1838
pyrethroid, 1838
target site resistance, 1838
olive fruit fly,
biological control, 1118
crude extract, 1118
microbial control, 1118
secondary metabolite, 1118
olive pyralid moth,
boring larva, 1602
branch protection zone, 1602
natural target pruning, 1602
pruning paint, 1602
Omatomidae,
Erotylidae, 305
Europs, 305
Loberus testaceus, 305
Nitidulidae, 305
one-choice,
habitat fidelity, 1932
spotted wing drosophila, 1932
4-way olfactometer, 1932
Ooencyrtus nezarae,
egg parasitism, 1973
Gryon japonicum, 1973
kairomone, 1973
soybean, 1973
OP-replacement insecticides,
biological control, 1126
generalist predator, 1126
green lacewing, 1126
reduced-risk insecticide, 1126
orange,
fumigation, 2341
lemon, 2341
phytosanitary, 2341
Vapormate, 2341
organophosphate,
Brevicoryne brassicae, 954
insecticide resistance, 954
neonicotinoid, 954
pyrethroid, 954
organosilicone,
almond, 1952
boscalid, 1952
diflubenzuron, 1952
pyraclostrobin, 1952
oriental cockroach,
Blatta orientalis, 2428
blattidae, 2428
invasive species, 2428
oriental fruit fly,
prezygotic compatibility, 695
sexual isolation, 695
ornamental nursery,
plant-based essential oil, 289
Pyrethroid, 289
Xylosandrus, 289
Oryzaephilus surinamensis,
pecan, 1071
Plodia interpunctella, 1071
stored-product pests, 1071
Oryza sativa, 837, 2585
biological control, 2585
Bt rice, 2585
cry gene, 2585
feeding guild, 837
injury equivalency, 837
rice, 837
rice planthopper, 837
Ostrinia furnacalis,
Cry1Ab/Cry2Aj protein, 2548
Cry1Ab/Vip3DA protein, 2548
insect resistance evaluation, 2548
transgenic maize, 2548
Ostrinia nubilalis,
distribution, 2084
pheromone, 2084
predictive model, 2084
spatial, 2084
ovarian development,
Apolygus lucorum, 1200
diet, 1200
follicle, 1200
Helicoverpa armigera egg, 1200
overwintering, 1574, 1619
corn leafhopper, 1574
incidence, 1574
mollicutes, 1574
northern corn rootworm, 1619
pest, 1574
population variation, 1619
overwintering adult,
brown marmorated stink bug, 2141
maximum temperature, 2141
population density, 2141
oviposition,
blueberry, 1767
cultivar, 1767
Drosophila suzukii, 1767
fruit firmness, 1767

P

Panonychus ulmi,
apple, 2118
development, 2118
life table, 2118
reproduction, 2118
Paralobesia viteana,
economic injury level, 905
grape, 905
integrated pest management (IPM), 905
parasitism,
biological control, 73
Eulophidae, 73
Platygastridae, 73
Vaccinium, 73
parasitoid behavior,
biological control, 1638
host-feeding, 1638
integrated pest management (IPM), 1638
life table, 1638
parasitoid community,
biological control, 1134
cabbage, 1134
insect–plant interaction, 1134
kale, 1134
parasitoid release,
establishment, 1145
invasive, 1145
natural enemy introduction, 1145
wood borers, 1145
pathogen detection,
Bactericera cockerelli, 1440
Candidatus Liberibacter solanacearum, 1440
LAMP, 1440
Zebra chip, 1440
PCR-restriction fragment-length polymorphism, 661, 2541
ITS, 661
mating assessment, 2541
Mediterranean fruit fly, 2541

ribosomal DNA, 661
species identification, 661
stored sperm identification, 2541
Tetranychus, 661
pea aphid,
Acyrthosiphon pisum, 2613
binomial sequential sampling, 2613
dry pea, 2613
Pisum sativum, 2613
peach fruit fly, 1608, 2336
cold treatment, 1608, 2336
Mediterranean fruit fly, 2336
phytosanitary treatment, 1608, 2336
phytosanitation, 1608
quarantine treatment, 1608, 2336
peanut, 587, 1112
entomopathogenic nematode, 1112
Frankliniella fusca, 587
Heterorhabditis, 1112
resistant genotypes, 587
Steinernema, 1112
Tospovirus, 587
white grub, 1112
pear psylla,
biological control, 776
generalist predator, 776
lethal, 776
reduced-risk insecticide, 776
sublethal effect, 776
pecan,
Oryzaephilus surinamensis, 1071
Plodia interpunctella, 1071
stored-product pests, 1071
Pectinophora gossypiella,
cryopreservation, 855
embryo, 855
permeabilization, 855
pink bollworm moth, 855
per capita reproduction,
biological control, 1979
Cleruchoides noackae, 1979
Mymaridae, 1979
Thaumastocoris peregrinus, 1979
Performance-2 model,
bootstrap percentile, 107
lower developmental threshold, 107
sum of effective temperatures, 107
upper developmental threshold, 107
Periplaneta japonica,
barcode, 2275
cockroach, 2275
invasive, 2275
pest, 2275
permeabilization,
cryopreservation, 855
embryo, 855
Pectinophora gossypiella, 855
pink bollworm moth, 855
permethrin,
bifenthrin, 329
Coptotermes acinaciformis, 329
Coptotermes formosanus, 329
pyrethroid, 329
peroxidase,
Blissus occiduus, 2568
Buchloë dactyloides, 2568
catalase, 2568
plant resistance, 2568
Persea borbonia,
laurel wilt, 1693
Raffaelea lauricola, 1693
redbay ambrosia beetle, 1693
pest, 1574, 2275
barcode, 2275
cockroach, 2275
corn leafhopper, 1574
incidence, 1574
invasive, 2275
mollicutes, 1574
overwintering, 1574
Periplaneta japonica, 2275
pesticide, 136, 150, 552, 756
agro-ecosystems, 756
behavior, 552
Bombus, 552
brown marmorated stink bug, 150
invasive species, 150
invertebrate, 756
mortality, 756
movement, 150
nontarget, 756
pollinator, 552
queen-less, 552
safety evaluation, 136
sublethal, 150
T. nubilale, 136
pesticide alternative, 64, 1986
Beauveria bassiana, 64, 1986
biological control, 64, 1986
resistance management, 64, 1986
pesticide mixture,
Diabrotica barberi, 1941
Diabrotica virgifera virgifera, 1941
pyramid, 1941
resistance management, 1941
pesticide toxicity,
chlorpyrifos, 1226
spinosad, 1226
sterile insect technique, 1226
Vienna-8, 1226
pest management, 247, 1726
antagonism, 247
bioassay, 247
broccoli, 1726
cauliflower, 1726
diel activity, 1726
horticulture, 247
stink bug, 1726
synergism, 247
pest monitoring,
cotton boll damage, 1310
plant bug, 1310
sucking bugs, 1310
Phenacoccus peruvianus,
phenology, 1486
sampling, 1486
spatial distribution, 1486
urban landscape, 1486
phenological indicator,
ambrosia beetle, 1780
bottle trap, 1780
forecasting activity, 1780
trap tree, 1780
phenology, 1486, 2491
Colorado potato beetle, 2491
diapause, 2491
insecticide resistance, 2491
integrated pest management (IPM), 2491
Phenacoccus peruvianus, 1486
sampling, 1486
spatial distribution, 1486
urban landscape, 1486
phenology model,
bronze beetle, 899
Chrysomelidae, 899
degree-day, 899
temperature, 899
pheromone, 482, 2084
distribution, 2084
grape, 482
mealybug, 482
monitoring, 482
Ostrinia nubilalis, 2084
predictive model, 2084
spatial, 2084
Washington state vineyard, 482
pheromone-baited traps,
clover, 844
Coleophora deauratella, 844
monitoring, 844
pheromone trap, 499, 1250
Bacillus thuringiensis, 1250
fumigation, 499
Helicoverpa zea, 1250
monitoring, 499
red flour beetle, 499
sulfuryl fluoride, 499
transgenic corn, 1250

Phloeomyzus passerinii,
cortical parenchyma, 473
Populus x canadensis, 473
tree resistance, 473
phosphine,
light brown apple moth, 1613
phytosanitary, 1613
postharvest pest control, 1613
quarantine treatment, 1613
phosphine resistance management,
insecticide resistance, 2208
lesser grain borer, 2208
red flour beetle, 2208
stored product, 2208
photoreceptor,
circadian clock, 450
clock genes, 450
cryptochrome, 450
Helicoverpa armigera, 450
phoxim,
Bt cotton, 375
emamectin benzoate, 375
fenvalerate, 375
Helicoverpa armigera, 375
Phthorimaea operculella, 1018, 1025
Bacillus thuringiensis, 1025
Bt, 1025
digestive enzymes, 1018
nutritional index, 1018
potato germplasms, 1018
Solanum tuberosum, 1025
transgenic, 1025
Phyllotreta spp.,
Crambe spp., 2557
flea beetle, 2557
glucosinolate, 2557
Sinapis alba, 2557
phylogenetic,
alfalfa weevil, 426
Hypera postica, 426
phylogenetic analysis,
COI, 408
genetic distance, 408
thrip, 408
phylogenetic relationships,
Liposcelis, 419
molecular identification, 419
morphological characteristics, 419
SEM, 419
physic nut,
Atta sexdens rubropilosa, 742
castor oil, 742
leaf-cutting ant, 742
seed lipid, 742
physiological indices,
apple cultivars, 1011
autumn shoots, 1011
Eriosoma lanigerum (Hausmann), 1011
resistance to insect, 1011
phytoplasma-insect relationships,
Auchenorrhyncha, 604
feeding-medium assay, 604
grapevine yellows, 604
phytoplasma seasonal variation, 604
phytoplasma seasonal variation,
Auchenorrhyncha, 604
feeding-medium assay, 604
grapevine yellows, 604
phytoplasma-insect relationships, 604
phytosanitary, 1179, 1613, 2267, 2341
biosecurity, 1179
fumigation, 2267, 2341
insect, 1179
lemon, 2341
light brown apple moth, 1613
mite, 1179
nitric oxide, 2267
orange, 2341
phosphine, 1613
plant quarantine, 1179
postharvest pest control, 1613, 2267
quarantine, 2267
quarantine treatment, 1613
Vapormate, 2341
phytosanitary treatment, 525, 1608, 2006, 2035, 2336
ATP, 2006
biochemical viability assay, 2006
cold treatment, 1608, 2336
colorimetric, 2006
ionizing radiation, 525
irradiation, 2035
Mediterranean fruit fly, 2336
peach fruit fly, 1608, 2336
phytosanitation, 525, 1608
postharvest, 2035
quarantine, 2006, 2035
quarantine pest, 525
quarantine treatment, 1608, 2336
systems approach, 2035
phytosanitation, 525, 1608, 2093
avocado, 2093
cold treatment, 1608
ionizing radiation, 525
peach fruit fly, 1608
phytosanitary treatment, 525, 1608
quarantine pest, 525
quarantine treatment, 1608
Raffaelea lauricola, 2093
redbay, 2093
Xyleborus glabratus, 2093
Phytoseiidae,
biological control, 1996
Galendromus occidentalis, 1996
inundative release, 1996
Tetranychidae, 1996
phytoseiids,
biological control, 80
Eotetranychus lewisi, 80
strawberry, 80
T. urticae, 80
phytovirus,
Aphididae, 683
electrical penetration graph, 683
host plant suitability, 683
Miscanthus sacchariflorus, 683
pink bollworm moth,
cryopreservation, 855
embryo, 855
Pectinophora gossypiella, 855
permeabilization, 855
Pinus strobus,
eastern white pine, 2391
Matsucoccidae, 2391
Matsucoccus macrocicatrices, 2391
southeastern United States, 2391
pirimicarb,
cyfluthrin, 400
imidacloprid, 400
metamidophos, 400
Myzus persicae, 400
pirimiphos methyl,
abamectin, 1848
acrinathrin, 1848
bifenazate, 1848
fenazaquin, 1848
Pisum sativum,
Acyrthosiphon pisum, 2613
binomial sequential sampling, 2613
dry pea, 2613
pea aphid, 2613
pitfall,
bed bug, 1802
carbon dioxide, 1802
chemical lure, 1802
sugar, 1802
yeast fermentation, 1802
plant-based essential oil,
ornamental nursery, 289
Pyrethroid, 289
Xylosandrus, 289
plant bug, 1310, 2077
cotton boll damage, 1310
cotton boll injury, 2077
Gossypium hirsutum, 2077
insect–plant interaction, 2077
pest monitoring, 1310
sucking bugs, 1310
plant community,
dispersal, 2296
Hemiptera vector, 2296
vineyard agroecosystem, 2296

plant compensation,
cotton, 1209
insect damage, 1209
lint yield, 1209
Lygus hesperus, 1209
plant development,
European corn borer, 1234
maize, 1234
Mediterranean corn borer, 1234
resistance, 1234
plant quarantine,
biosecurity, 1179
insect, 1179
mite, 1179
phytosanitary, 1179
plant resistance, 1029, 1045, 1456, 2568
Blissus occiduus, 2568
Buchloë dactyloides, 2568
catalase, 2568
cotton fleahopper, 1045
cross-resistance, 1029
Cylas formicarius, 1456
Diabrotica, 1456
Gossypium hirsutum, 1045
Ipomoea, 1456
native species, 1029
peroxidase, 2568
Pseudatomoscelis seriatus, 1045
wheat pests, 1029
white grub, 1456
plant stress,
arthropod predators, 1260
biological control, 1260
cotton, 1260
whitefly, 1260
plant virus,
acquisition, 1878
life history, 1878
resistance, 1878
transmission, 1878
plastic cable sheathings,
Coptotermes, 1395
nylon 12, 1395
polyethylene, 1395
subterranean termite, 1395
Platygastridae,
biological control, 73
Eulophidae, 73
parasitism, 73
Vaccinium, 73
Plodia interpunctella, 1071, 1503
economic analysis, 1503
esfenvalerate, 1503
methoprene, 1503
Oryzaephilus surinamensis, 1071
pecan, 1071
stored-product pests, 1071
Plutella xylostella, 1075, 1103, 1423
additives, 1075
Bacillus thuringiensis, 1075
baseline toxicity, 1423
bioassay, 1075
biocontrol, 1075
broccoli, 1103
Diadegma insulare, 1103
diamondback moth, 1103
indoxacarb, 1423
metaflumizone, 1423
simulation, 1103
pollen diets,
fat body, 1553
honey bee, 1553
nutritional status, 1553
protein, 1553
pollen germination,
Echinacea angustifolia, 2055
insect pollinator, 2055
pollen-tube quantification, 2055
shrivelled (retracted) styles, 2055
pollen-tube quantification,
Echinacea angustifolia, 2055
insect pollinator, 2055
pollen germination, 2055
shrivelled (retracted) styles, 2055
pollination, 543, 576
alfalfa, 543
alfalfa-seed production, 576
Medicago sativa, 576
Megachile rotundata, 543, 576
Megachilidae, 543, 576
reproduction, 543
pollinator,
behavior, 552
Bombus, 552
pesticide, 552
queen-less, 552
polyandry,
Helicoverpa armigera, 708
mating history, 708
sperm competition, 708
sperm transfer, 708
polyethylene,
Coptotermes, 1395
nylon 12, 1395
plastic cable sheathings, 1395
subterranean termite, 1395
polymer matrix,
methyl eugenol, 727
monitoring traps, 727
spinosad, 727
toxicant, 727
polytene chromosome,
Bactrocera cucurbitae, 995
Bactrocera dorsalis, 995
genetic sexing strain, 995
Tephritidae, 995
Popillia japonica, 1190, 2048
corn, 2048
harmonization plan, 1190
insecticide, 1190
Japanese beetle, 2048
neonicotinoid, 1190
scarab, 1190
silk clipping, 2048
population,
Bt rice, 1887
detoxification enzyme, 1887
ecological fitness, 1887
Nilaparvata lugens, 1887
population density,
brown marmorated stink bug, 2141
maximum temperature, 2141
overwintering adult, 2141
population genetic, 382, 1355
Bemisia tabaci, 1355
insect, 382
invasive species, 1355
stochastic processes, 382
population parameter,
bio-insecticide, 2177
finite rate of increase, 2177
intrinsic rate of increase, 2177
net reproductive rate, 2177
population projection,
bootstrap, 1
Chrysopa pallens, 1
jackknife, 1
life table, 1
population structure,
AFLP, 2225
cotton pest, 2225
dispersal, 2225
gene flow, 2225
population variation,
northern corn rootworm, 1619
overwintering, 1619
Populus x canadensis,
cortical parenchyma, 473
Phloeomyzus passerinii, 473
tree resistance, 473
postharvest,
irradiation, 2035
phytosanitary treatment, 2035
quarantine, 2035
systems approach, 2035
postharvest fumigation,
Epiphyas postvittana, 1155
methyl bromide, 1155
sorption kinetics, 1155
postharvest pest control, 1613, 2267
fumigation, 2267
light brown apple moth, 1613
nitric oxide, 2267
phosphine, 1613
phytosanitary, 1613, 2267
quarantine, 2267
quarantine treatment, 1613

potato, 786, 1964
acquisition access period, 1964
damage, 786
Liberibacter, 1964
mortality, 786
potato psyllid, 1964
recovery, 786
tomato, 1964
wireworm, 786
potato germplasms,
digestive enzymes, 1018
nutritional index, 1018
Phthorimaea operculella, 1018
potato psyllid,
acquisition access period, 1964
Liberibacter, 1964
potato, 1964
tomato, 1964
potyviruses,
aphid, 1566
morning glory, 1566
sweet potato, 1566
predatory mite,
litchi, 2183
litchi erineum mite, 2183
selectivity, 2183
predictive model,
distribution, 2084
Ostrinia nubilalis, 2084
pheromone, 2084
spatial, 2084
preference,
host-plant resistance, 2577
nutritional index, 2577
Rag1 gene, 2577
soybean, 2577
presence–absence,
full-count, 1707
mint, 1707
twospotted spider mite, 1707
prey consumption,
adult longevity, 2621
Cheilomenes sexmaculata development, 2621
fertility, 2621
prezygotic compatibility,
oriental fruit fly, 695
sexual isolation, 695
principal component analysis,
multivariate statistical analysis, 1053
tritrophic interaction, 1053
VOCs, 1053
prism trap,
detection tools, 2407
multifunnel trap, 2407
Sabic green, 2407
Sabic purple, 2407
probability density function,
Coptotermes formosanus, 2442
Isoptera, 2442
mark–recapture protocol, 2442
Rhinotermitidae, 2442
probiotic,
irradiation, 641
medfly, 641
sperm, 641
sterile insect technique, 641
production,
insect colony effect, 50
SeMNPV, 50
Spexit, 50
Vir-ex, 50
protein,
fat body, 1553
honey bee, 1553
nutritional status, 1553
pollen diets, 1553
proteinases,
Cry1Ac resistance, 2514
Helicoverpa armigera, 2514
nonbinding site, 2514
protein expression,
CYP4, 393
cytochrome P450 monooxygenase, 393
Diaphorina citri, 393
gene expression, 393
pruning paint,
boring larva, 1602
branch protection zone, 1602
natural target pruning, 1602
olive pyralid moth, 1602
Prunus dulcis,
Amyelois transitella, 1365
bird damage, 1365
Corvidae, 1365
degree-day, 1365
survivorship, 1365
Pseudatomoscelis seriatus,
cotton fleahopper, 1045
Gossypium hirsutum, 1045
plant resistance, 1045
Pseudomonas aeruginosa,
carcass-burying behavior, 1812
Coptotermes formosanus, 1812
disease resistance, 1812
lufenuron, 1812
Psocoptera,
booklouse, 491
insect pest, 491
Liposcelis, 491
stored product, 491
Psyllidae,
genetic differentiation, 1004
Homoptera, 1004
ISSR, 1004
molecular marker, 1004
puparial survival,
hay drying, 1164
Mayetiola destructor (Say), 1164
quarantine control, 1164
systems approach, 1164
purple,
Agrilus planipennis, 2415
fluon, 2415
green, 2415
multifunnel trap, 2415
pyraclostrobin,
almond, 1952
boscalid, 1952
diflubenzuron, 1952
organosilicone, 1952
pyramid,
Diabrotica barberi, 1941
Diabrotica virgifera virgifera, 1941
pesticide mixture, 1941
resistance management, 1941
pyrethroid, 289, 329, 954, 959, 1838
Aedes aegypti, 959
bifenthrin, 329
Brevicoryne brassicae, 954
carbamate, 1838
Coptotermes acinaciformis, 329
Coptotermes formosanus, 329
cross-resistance, 959
insecticide resistance, 954, 959
Myzus persicae, 1838
neonicotinoid, 954
oilseed rape, 1838
organophosphate, 954
ornamental nursery, 289
permethrin, 329
plant-based essential oil, 289
target site resistance, 1838
Xylosandrus, 289
pyrethroid insecticide resistance,
Culex pipiens pallens, 360
knockdown resistance, 360
susceptibility, 360

Q

quantitative PCR,
Bemisia tabaci, 1446
endogenous reference gene, 1446
gene expression, 1446
RNA-seq, 1446
quarantine, 520, 2006, 2020, 2035, 2267
apple, 520
ATP, 2006
biochemical viability assay, 2006
colorimetric, 2006
diapause, 520
export, 520
fumigation, 2267
generic dose, 2020
irradiation, 2020, 2035
Mediterranean fruit fly, 2020
modified atmosphere, 2020
nitric oxide, 2267

phytosanitary, 2267
phytosanitary treatment, 2006, 2035
postharvest, 2035
postharvest pest control, 2267
systems approach, 2035
quarantine control,
hay drying, 1164
Mayetiola destructor (Say), 1164
puparial survival, 1164
systems approach, 1164
quarantine pest,
ionizing radiation, 525
phytosanitary treatment, 525
phytosanitation, 525
quarantine treatment, 1608, 1613, 2336
cold treatment, 1608, 2336
light brown apple moth, 1613
Mediterranean fruit fly, 2336
peach fruit fly, 1608, 2336
phosphine, 1613
phytosanitary, 1613
phytosanitary treatment, 1608, 2336
phytosanitation, 1608
postharvest pest control, 1613
queen-less,
behavior, 552
Bombus, 552
pesticide, 552
pollinator, 552
Quercus virginiana,
alternating generation, 1747
bullet gall, 1747
gall wasp, 1747

R

Raffaelea lauricola, 1693, 2093
avocado, 2093
laurel wilt, 1693
Persea borbonia, 1693
phytosanitation, 2093
redbay, 2093
redbay ambrosia beetle, 1693
Xyleborus glabratus, 2093
Rag1 gene,
host-plant resistance, 2577
nutritional index, 2577
preference, 2577
soybean, 2577
(R)-carvone,
Epimedium pubescens, 513
Liposcelis bostrychophila, 513
repellency, 513
Tribolium castaneum, 513
Rdl gene,
cross-resistance, 2167
dieldrin, 2167
fipronil, 2167
insecticide resistance, 2167
rearing,
first-generation, 533
life table, 533
Megacopta cribraria, 533
soybean, 533
receptor binding,
Aedes albopictus, 1098
Bacillus thuringiensis, 1098
ELISA, 1098
immunohistochemistry, 1098
recovery,
damage, 786
mortality, 786
potato, 786
wireworm, 786
redbay,
avocado, 2093
phytosanitation, 2093
Raffaelea lauricola, 2093
Xyleborus glabratus, 2093
redbay ambrosia beetle, 1693, 2286
chemical control, 2286
exotic insect, 2286
insect–disease complex, 2286
Lauraceae, 2286
laurel wilt, 1693
Persea borbonia, 1693
Raffaelea lauricola, 1693
red flour beetle, 499, 2208
fumigation, 499
insecticide resistance, 2208
lesser grain borer, 2208
monitoring, 499
pheromone trap, 499
phosphine resistance management, 2208
stored product, 2208
sulfuryl fluoride, 499
red palm weevil,
ethyl acetate, 1739
integrated pest management (IPM), 1739
kairomone, 1739
trap, 1739
reduced-risk insecticide, 776, 1126
biological control, 776, 1126
generalist predator, 776, 1126
green lacewing, 1126
lethal, 776
OP-replacement insecticides, 1126
pear psylla, 776
sublethal effect, 776
refuge, 816, 1525
Bacillus thuringiensis, 1525
compensation, 816
corn, 816
Diatraea grandiosella, 816
field-evolved resistance, 1525
insect resistance management, 1525
southwestern corn borer, 816
refuge strategy,
Diabrotica virgifera virgifera, 2195
event 59122, 2195
resistance management, 2195
regression analysis,
Chilo suppressalis, 1832
dynamics, 1832
insecticide resistance, 1832
repellency,
Epimedium pubescens, 513
Liposcelis bostrychophila, 513
(R)-carvone, 513
Tribolium castaneum, 513
repellent, 131, 2522
bed bug, 2522
bioactive compound, 131
clove oil, 131
DEET, 2522
essential oil, 2522
natural product, 2522
Solenopsis invicta, 131
toxicity, 131
repellent netting,
alphacypermethrin, 1699
Aphidius colemani, 1699
cabbage, 1699
Myzus persicae, 1699
reproduction, 543, 2118
alfalfa, 543
apple, 2118
development, 2118
life table, 2118
Megachile rotundata, 543
Megachilidae, 543
Panonychus ulmi, 2118
pollination, 543
reproductive performance,
Bt toxin, 614
Helicoverpa armigera, 614
lifespan, 614
Spodoptera exigua, 614
resistance, 1234, 1414, 1430, 1473, 1878, 2530, 2602
acquisition, 1878
antixenosis, 2602
Bemisia tabaci, 1414
Brassica oleracea, 2602
cotton aphid, 1430
English grain aphid, 1473
European corn borer, 1234
flonicamid, 1430
germplasm, 1473
host plant, 1414
IGRs, 1414
insecticide formulation, 2530
insecticide susceptibility, 2530
insecticide toxicity, 2530
life history, 1878

light reflectance, 2602
maize, 1234
Mediterranean corn borer, 1234
neonicotinoid, 1414, 1430
plant development, 1234
plant virus, 1878
sulfoxaflor, 1430
Thrips tabaci, 2602
tolerance, 1473
transmission, 1878
wheat, 1473
resistance management, 64, 823, 1941, 1986, 2195
Beauveria bassiana, 64, 1986
biological control, 64, 1986
Diabrotica barberi, 1941
Diabrotica virgifera virgifera, 823, 1941, 2195
event 59122, 2195
Event MON 88017, 823
Event MON 89034, 823
pesticide alternative, 64, 1986
pesticide mixture, 1941
pyramid, 1941
refuge strategy, 2195
transgenic corn, 823
resistance monitoring, 945, 1855, 2190
baseline susceptibility, 2190
beet armyworm, 1855
Chilo suppressalis, 2190
chlorantraniliprole, 2190
gel bait, 945
German cockroach, 945
indoxacarb, 945
insecticide resistance, 945, 1855
resistance ratio, 2190
resistance ratio,
baseline susceptibility, 2190
Chilo suppressalis, 2190
chlorantraniliprole, 2190
resistance monitoring, 2190
resistance to insect,
apple cultivars, 1011
autumn shoots, 1011
Eriosoma lanigerum (Hausmann), 1011
physiological indices, 1011
resistant genotypes,
Frankliniella fusca, 587
peanut, 587
Tospovirus, 587
respiration,
diapause, 1089
Megachile rotundata, 1089
metabolic rate, 1089
thermal stress, 1089
Reticulitermes,
fipronil, 2160
foraging behavior, 2160
termite, 2160
termiticide, 2160
Rhagoletis indifferens,
apple maggot, 1511
Malus spectabilis, 1511
microsatellites, 1511
western cherry fruit fly, 1511
Rhinotermitidae,
Coptotermes formosanus, 2442
Isoptera, 2442
mark–recapture protocol, 2442
probability density function, 2442
Rhopalosiphum padi,
Schizaphis graminum, 1894
Sitobion avenae, 1894
specific resistance, 1894
trait hierarchy, 1894
ribosomal DNA,
ITS, 661
PCR-restriction fragment-length polymorphism, 661
species identification, 661
Tetranychus, 661
rice,
feeding guild, 837
injury equivalency, 837
Oryza sativa, 837
rice planthopper, 837
rice planthopper,
feeding guild, 837
injury equivalency, 837
Oryza sativa, 837
rice, 837
rice water weevil,
chlorantraniliprole, 181
seed treatment, 181
sublethal effect, 181
thiamethoxam, 181
ripeness,
cranberry, 2424
fruit, 2424
spotted wing drosophila, 2424
susceptibility, 2424
RNA-seq,
Bemisia tabaci, 1446
endogenous reference gene, 1446
gene expression, 1446
quantitative PCR, 1446
royal jelly,
bee strain, 1958
botanical origin, 1958
geographical origin, 1958
10-hydroxy-*trans*-2-decenoic acid, 1958
Russian honey bee,
grooming behavior, 566
mite fall, 566
Varroa destructor, 566
Varroa resistance, 566
Russian wheat aphid,
alternative host plant, 2595
biotype, 2595
Diuraphis noxia, 2595
intrinsic rate of increase, 2595

S

Sabic green,
detection tools, 2407
multifunnel trap, 2407
prism trap, 2407
Sabic purple, 2407
Sabic purple,
detection tools, 2407
multifunnel trap, 2407
prism trap, 2407
Sabic green, 2407
safe pesticide,
Diglyphus isaea, 1590
greenhouse biocontrol, 1590
greenhouse pest management, 1590
leafminer biocontrol, 1590
safety evaluation,
pesticide, 136
T. nubilale, 136
Saffron Thread Yellow,
Sunny Summer Yellow, 2109
Tartar Red, 2109
trap, 2109
western cherry fruit fly, 2109
salvage logging,
CT scan, 1331
damage progression, 1331
fire-killed trees, 1331
whitespotted sawyer, 1331
sampling,
Phenacoccus peruvianus, 1486
phenology, 1486
spatial distribution, 1486
urban landscape, 1486
scarab,
harmonization plan, 1190
insecticide, 1190
neonicotinoid, 1190
Popillia japonica, 1190
scarab larvae,
botanical insecticides, 159
essential oil, 159
GC-MS, 159
white grub, 159
Schizaphis graminum,
Rhopalosiphum padi, 1894
Sitobion avenae, 1894
specific resistance, 1894
trait hierarchy, 1894
Scolytinae,
Cerambycidae, 206
Cleridae, 206
Histeridae, 206
Trogossitidae, 206

screening protocols,
honey bee, 1535
surveillance, 1535
Tropilaelaps mite, 1535
SDS-PAGE,
Bacillus thuringiensis, 36
gamma radiation, 36
kilogray, 36
toxicity, 36
seasonal effect,
Apis mellifera, 2280
food availability, 2280
honey bee, 2280
hygienic behavior, 2280
secondary metabolite,
biological control, 1118
crude extract, 1118
microbial control, 1118
olive fruit fly, 1118
secondary screwworm,
carrion, 1927
forensic entomology, 1927
myiasis, 1927
spray-dried blood, 1927
seedling damage,
dead-heart plants, 1669
Diabrotica undecimpunctata howardi, 1669
Maize, 1669
southern corn rootworm, 1669
seed lipid,
Atta sexdens rubropilosa, 742
castor oil, 742
leaf-cutting ant, 742
physic nut, 742
seed treatment, 181, 2384
chlorantraniliprole, 181
electronic penetration graph, 2384
rice water weevil, 181
soybean aphid, 2384
sublethal effect, 181
thiamethoxam, 181, 2384
seed yield,
Allium cepa L., 1386
floral traits, 1386
male sterility, 1386
nectar composition, 1386
selective insecticide,
integrated pest management (IPM), 2126
systemics, 2126
water volume, 2126
selectivity,
litchi, 2183
litchi erineum mite, 2183
predatory mite, 2183
SEM,
Liposcelis, 419
molecular identification, 419
morphological characteristics, 419
phylogenetic relationships, 419
seminatural areas pollinator habitat,
agricultural landscape, 97
crop pollination, 97
wild bees, 97
semiochemical,
antiaggregant, 221
lodgepole pine, 221
mountain pine beetle, 221
verbenone, 221
semipersistent,
leaf disk transmission assay, 16
vector, 16
SeMNPV,
insect colony effect, 50
production, 50
Spexit, 50
Vir-ex, 50
Sesamia nonagrioides,
borer damage, 2241
fumonisin, 2241
Fusarium verticillioides, 2241
Zea mays, 2241
sex pheromone,
flight tunnel, 2144
insect behavior, 2144
λ-cyhalothrin, 2144
lethal effect, 2144
sex-specific mortality,
bed bug, 988
Cimex lectularius, 988
deltamethrin, 988
sexual behavior,
Anastrepha ludens, 735
genetic sexing, 735
sterile insect technique, 735
Tapachula-7, 735
sexual isolation,
oriental fruit fly, 695
prezygotic compatibility, 695
sexual maturation,
adult starvation, 215
daily reproductive rhythm, 215
Glenea cantor, 215
lifetime reproductive pattern, 215
sexual selection,
courtship, 1373
mating behavioral sequence, 1373
mating success, 1373
shrivelled (retracted) styles,
Echinacea angustifolia, 2055
insect pollinator, 2055
pollen germination, 2055
pollen-tube quantification, 2055
side-shoot,
apical-meristem injury, 1339
cranberry, 1339
fruit set, 1339
tipworm, 1339
silica gel,
essential oil, 1902
Juniperus oxycedrus ssp. *oxycedrus*, 1902
Sitophilus oryzae, 1902
stored wheat, 1902
Tribolium confusum, 1902
silk clipping,
corn, 2048
Japanese beetle, 2048
Popillia japonica, 2048
simulation, 1103, 2473
broccoli, 1103
Callosobruchus maculatus, 2473
cowpea, 2473
Diadegma insulare, 1103
diamondback moth, 1103
insect resistance management, 2473
Plutella xylostella, 1103
transgenic crop, 2473
Sinapis alba,
Crambe spp., 2557
flea beetle, 2557
glucosinolate, 2557
Phyllotreta spp., 2557
Sitobion avenae,
Rhopalosiphum padi, 1894
Schizaphis graminum, 1894
specific resistance, 1894
trait hierarchy, 1894
Sitodiplosis mosellana,
biotic factor, 2043
environmental factor, 2043
flight ability, 2043
flight mill, 2043
Sitophilus granarius,
insect infestation, 1058
mineral, 1058
storage, 1058
vitamin, 1058
Sitophilus oryzae,
essential oil, 1902
Juniperus oxycedrus ssp. *oxycedrus*, 1902
silica gel, 1902
stored wheat, 1902
Tribolium confusum, 1902
six-spotted thrip,
action threshold, 1757
degree-day, 1757
Fragaria ananassa, 1757
mite-day, 1757
size restriction,
Cicadellidae, 2027
Dracaena marginata, 2027
nursery stock, 2027

offshore preclearance program, 2027
sodium borate,
ant, 929
bait, 929
boric acid, 929
feeding, 929
soil-applied insecticides,
acephate, 807
aldicarb, 807
imidacloprid, 807
thiamethoxam, 807
Solanum tuberosum,
Bacillus thuringiensis, 1025
Bt, 1025
Phthorimaea operculella, 1025
transgenic, 1025
solarization,
debarking, 235
firewood, 235
grinding, 235
oak mortality, 235
Solenopsis invicta,
bioactive compound, 131
clove oil, 131
repellent, 131
toxicity, 131
soluble sugar content,
fruit safety, 299
fruit shelf life, 299
Mangifera indica, 299
Oecophylla smaragdina, 299
sorption kinetics,
Epiphyas postvittana, 1155
methyl bromide, 1155
postharvest fumigation, 1155
South American fruit fly,
azadirachtin, 1772
Mediterranean fruit fly, 1772
neem, 1772
toxic bait, 1772
southeastern United States,
eastern white pine, 2391
Matsucoccidae, 2391
Matsucoccus macrocicatrices, 2391
Pinus strobus, 2391
southern corn rootworm,
dead-heart plants, 1669
Diabrotica undecimpunctata howardi, 1669
Maize, 1669
seedling damage, 1669
southwestern corn borer,
compensation, 816
corn, 816
Diatraea grandiosella, 816
refuge, 816
soybean, 533, 1317, 1479, 1676, 1973, 2577
Aphis glycines, 1479
biotype, 1479
egg parasitism, 1973
first-generation, 533
Gryon japonicum, 1973
Halyomorpha halys, 1317
host-plant resistance, 1479, 2577
invasive species, 1676
kairomone, 1973
life table, 533
Megacopta cribraria, 533, 1676
native stink bugs, 1317
nutritional index, 2577
Ooencyrtus nezarae, 1973
preference, 2577
Rag1 gene, 2577
rearing, 533
thresholds, 1317
soybean aphid,
electronic penetration graph, 2384
seed treatment, 2384
thiamethoxam, 2384
soybean genotype,
aphid biotype, 2234
EPG, 2234
feeding behavior, 2234
spatial,
distribution, 2084
Ostrinia nubilalis, 2084
pheromone, 2084
predictive model, 2084
spatial distribution,
Phenacoccus peruvianus, 1486
phenology, 1486
sampling, 1486
urban landscape, 1486
species distribution model,
invasive species, 311
subterranean termite, 311
urban pest, 311
species identification,
ITS, 661
PCR-restriction fragment-length polymorphism, 661
ribosomal DNA, 661
Tetranychus, 661
specific resistance,
Rhopalosiphum padi, 1894
Schizaphis graminum, 1894
Sitobion avenae, 1894
trait hierarchy, 1894
sperm,
irradiation, 641
medfly, 641
probiotic, 641
sterile insect technique, 641
sperm competition,
Helicoverpa armigera, 708
mating history, 708
polyandry, 708
sperm transfer, 708
sperm transfer,
Helicoverpa armigera, 708
mating history, 708
polyandry, 708
sperm competition, 708
Spexit,
insect colony effect, 50
production, 50
SeMNPV, 50
Vir-ex, 50
Sphaerophoria spp.,
conservation biological control, 2327
flowering plant, 2327
habitat management, 2327
natural enemy, 2327
spinosad, 727, 1226, 1825, 1911
acute toxicity, 1825
biological parameter, 1825
chlorpyrifos, 1226
chlorpyrifos-methyl plus deltamethrin, 1911
field strains, 1911
methyl eugenol, 727
monitoring traps, 727
mortality, 1911
pesticide toxicity, 1226
polymer matrix, 727
Spodoptera exigua, 1825
sterile insect technique, 1226
stored-grain insects, 1911
sublethal effect, 1825
toxicant, 727
Vienna-8, 1226
spirotetramat,
cyantraniliprole, 597
Cyazypyr, 597
fluorescein sodium salt, 597
imidacloprid, 597
spittlebug,
adult damage, 1871
Brachiaria, 1871
host-plant resistance, 1871
Spodoptera exigua, 614, 800, 1240, 1825
acute toxicity, 1825
biological parameter, 1825
Bt toxin, 614
Helicoverpa armigera, 614
insect-plant interaction, 1240
instar determination method, 800
kernel density estimation, 800
lifespan, 614
N availability, 1240
nitrogen, 800
reproductive performance, 614
spinosad, 1825
sublethal effect, 1825
Zea mays, 1240

spotted wing drosophila, 1932, 2424
cranberry, 2424
fruit, 2424
habitat fidelity, 1932
one-choice, 1932
ripeness, 2424
susceptibility, 2424
4-way olfactometer, 1932
spray-dried blood,
carrion, 1927
forensic entomology, 1927
myiasis, 1927
secondary screwworm, 1927
16S rDNA,
Cacopsylla chinensis, 536
cytochrome oxidase I, 536
statistical power,
detection capacity, 1659
environmental risk assessment, 1659
insect, 1659
transgenic crop, 1659
Steinernema, 622, 1112
Diabrotica barberi, 622
Diabrotica virgifera virgifera, 622
entomopathogenic nematode, 1112
Heterorhabditis, 622, 1112
Metarhizium, 622
peanut, 1112
white grub, 1112
Steinernema carpocapsae,
entomopathogenic nematode, 648
integrated pest management (IPM), 648
Steinernematidae, 648
Tenebrionidae, 648
Steinernematidae,
entomopathogenic nematode, 648
integrated pest management (IPM), 648
Steinernema carpocapsae, 648
Tenebrionidae, 648
sterile insect technique, 641, 735, 1226
Anastrepha ludens, 735
chlorpyrifos, 1226
genetic sexing, 735
irradiation, 641
medfly, 641
pesticide toxicity, 1226
probiotic, 641
sexual behavior, 735
sperm, 641
spinosad, 1226
Tapachula-7, 735
Vienna-8, 1226
stink bug, 912, 1726
blackberry, 912
broccoli, 1726
cauliflower, 1726
diel activity, 1726
Euschistus spp., 912
injury, 912
monitoring, 912
pest management, 1726
stochastic processes,
insect, 382
population genetic, 382
stone pine,
cone abortion, 229
Ernobius spp., 229
Leptoglossus occidentalis, 229
nut production, 229
storage,
insect infestation, 1058
mineral, 1058
Sitophilus granarius, 1058
vitamin, 1058
stored-grain insects,
chlorpyrifos-methyl plus deltamethrin, 1911
field strains, 1911
mortality, 1911
spinosad, 1911
stored product, 491, 2208
booklouse, 491
insecticide resistance, 2208
insect pest, 491
lesser grain borer, 2208
Liposcelis, 491
phosphine resistance management, 2208
Psocoptera, 491
red flour beetle, 2208
stored-product pests,
Oryzaephilus surinamensis, 1071
pecan, 1071
Plodia interpunctella, 1071
stored sperm identification,
mating assessment, 2541
Mediterranean fruit fly, 2541
PCR-restriction fragment-length polymorphism, 2541
stored wheat,
essential oil, 1902
Juniperus oxycedrus ssp. *oxycedrus*, 1902
silica gel, 1902
Sitophilus oryzae, 1902
Tribolium confusum, 1902
strawberry,
biological control, 80
Eotetranychus lewisi, 80
phytoseiids, 80
T. urticae, 80
Striacosta albicosta,
economic injury level, 1274
economic threshold, 1274
larval survival, 1274
western bean cutworm, 1274
sublethal,
brown marmorated stink bug, 150
invasive species, 150
movement, 150
pesticide, 150
sublethal effect, 10, 181, 776, 1825
acute toxicity, 1825
biological control, 776
biological parameter, 1825
chlorantraniliprole, 181
fecundity, 10
generalist predator, 776
lethal, 776
Nilaparvata lugens, 10
pear psylla, 776
reduced-risk insecticide, 776
rice water weevil, 181
seed treatment, 181
spinosad, 1825
Spodoptera exigua, 1825
survival, 10
thiamethoxam, 181
wing formation, 10
subterranean termite, 311, 1395, 1794
bait, 1794
Coptotermes, 1395
Coptotermes curvignathus, 1794
Coptotermes gestroi, 1794
invasive species, 311
nylon 12, 1395
plastic cable sheathings, 1395
polyethylene, 1395
species distribution model, 311
sugar, 1794
urban pest, 311
sucking bugs,
cotton boll damage, 1310
pest monitoring, 1310
plant bug, 1310
sugar, 1794, 1802
bait, 1794
bed bug, 1802
carbon dioxide, 1802
chemical lure, 1802
Coptotermes curvignathus, 1794
Coptotermes gestroi, 1794
pitfall, 1802
subterranean termite, 1794
yeast fermentation, 1802
sulfoxaflor,
cotton aphid, 1430
flonicamid, 1430
neonicotinoid, 1430
resistance, 1430

sulfuryl fluoride,
fumigation, 499
monitoring, 499
pheromone trap, 499
red flour beetle, 499
sum of effective temperatures,
bootstrap percentile, 107
lower developmental threshold, 107
Performance-2 model, 107
upper developmental threshold, 107
Sunny Summer Yellow,
Saffron Thread Yellow, 2109
Tartar Red, 2109
trap, 2109
western cherry fruit fly, 2109
supercooling point,
bed bug, 2433
cold tolerance, 2433
freezing, 2433
nonchemical control, 2433
surface treatment,
binary combinations, 1064
cyfluthrin, 1064
imidacloprid, 1064
surveillance,
honey bee, 1535
screening protocols, 1535
Tropilaelaps mite, 1535
survey,
degree-day, 1495
distribution, 1495
invasive species ecology, 1495
survival,
fecundity, 10
Nilaparvata lugens, 10
sublethal effect, 10
wing formation, 10
survival rate,
Bemisia tabaci gossypol, 1379
development, 1379
survivorship,
Amyelois transitella, 1365
bird damage, 1365
Corvidae, 1365
degree-day, 1365
Prunus dulcis, 1365
susceptibility, 360, 2424
cranberry, 2424
Culex pipiens pallens, 360
fruit, 2424
knockdown resistance, 360
pyrethroid insecticide resistance, 360
ripeness, 2424
spotted wing drosophila, 2424
sweet potato,
aphid, 1566
morning glory, 1566
potyviruses, 1566
synergism,
antagonism, 247
bioassay, 247
horticulture, 247
pest management, 247
synergist,
Bactrocera dorsalis, 970
carboxylesterases, 970
toxicity, 970
systemic insecticide,
canopy translocation, 2399
hemlock woolly adelgid, 2399
neonicotinoid, 2399
Tsuga canadensis, 2399
systemics,
integrated pest management (IPM), 2126
selective insecticide, 2126
water volume, 2126
systemic uptake bioassay,
desert environment, 1404
host plant, 1404
insecticide resistance, 1404
systems approach, 1164, 2035
hay drying, 1164
irradiation, 2035
Mayetiola destructor (Say), 1164
phytosanitary treatment, 2035
postharvest, 2035
puparial survival, 1164
quarantine, 2035
quarantine control, 1164

T

Tapachula-7,
Anastrepha ludens, 735
genetic sexing, 735
sexual behavior, 735
sterile insect technique, 735
target site resistance,
carbamate, 1838
Myzus persicae, 1838
oilseed rape, 1838
pyrethroid, 1838
Tartar Red,
Saffron Thread Yellow, 2109
Sunny Summer Yellow, 2109
trap, 2109
western cherry fruit fly, 2109
tau-fluvalinate,
amitraz, 558
honey bee, 558
miticides, 558
Varroa destructor, 558
temperature,
bronze beetle, 899
Chrysomelidae, 899
degree-day, 899
phenology model, 899
temperature-dependent development,
biophysical model, 124
degree-day, 124
western tarnished plant bug, 124
temperature effects,
acoustic emissions, 338
drywood termites, 338
feeding periodicity, 338
Tenebrionidae,
entomopathogenic nematode, 648
integrated pest management (IPM), 648
Steinernema carpocapsae, 648
Steinernematidae, 648
Tephritidae,
Bactrocera cucurbitae, 995
Bactrocera dorsalis, 995
genetic sexing strain, 995
polytene chromosome, 995
termite, 2160, 2216
breeding system, 2216
fipronil, 2160
foraging behavior, 2160
invasive species, 2216
neoteny, 2216
Reticulitermes, 2160
termiticide, 2160
termiticide,
fipronil, 2160
foraging behavior, 2160
Reticulitermes, 2160
termite, 2160
Tetranychidae,
biological control, 1996
Galendromus occidentalis, 1996
inundative release, 1996
Phytoseiidae, 1996
Tetranychus, 463, 661
cryptic species, 463
cytochrome *c* oxidase subunit I, 463
DNA-based identification, 463
internal transcribed spacer, 463
ITS, 661
PCR-restriction fragment-length polymorphism, 661
ribosomal DNA, 661
species identification, 661
Tetranychus cinnabarinus,
F1538I mutation, 2457
fenpropathrin, 2457
molecular marker, 2457
voltage-gate sodium channel, 2457

Tetranychus urticae,
cotton, 862
twospotted spider mite, 862
yield, 862
Thaumastocoris peregrinus,
biological control, 1979
Cleruchoides noackae, 1979
Mymaridae, 1979
per capita reproduction, 1979
thermal requirement,
mushroom, 115
nonlinear modeling, 115
variety, 115
thermal stress,
diapause, 1089
Megachile rotundata, 1089
metabolic rate, 1089
respiration, 1089
thiamethoxam, 181, 807, 1302, 2384
acephate, 807
aldicarb, 807
Aphis glycines, 1302
chlorantraniliprole, 181
electronic penetration graph, 2384
imidacloprid, 807
integrated pest management (IPM), 1302
rice water weevil, 181
seed treatment, 181, 2384
soil-applied insecticides, 807
soybean aphid, 2384
sublethal effect, 181
thresholds,
Halyomorpha halys, 1317
native stink bugs, 1317
soybean, 1317
thrip,
COI, 408
genetic distance, 408
phylogenetic analysis, 408
Thrips tabaci,
antixenosis, 2602
Brassica oleracea, 2602
light reflectance, 2602
resistance, 2602
tipworm,
apical-meristem injury, 1339
cranberry, 1339
fruit set, 1339
side-shoot, 1339
T. nubilale,
pesticide, 136
safety evaluation, 136
tobacco budworm,
Bt cotton, 1819
insect resistance management, 1819
tolerance, 1036, 1473
Aphis glycines, 1036
English grain aphid, 1473
germplasm, 1473
host-plant resistance, 1036
resistance, 1473
wheat, 1473
tomato,
acquisition access period, 1964
Liberibacter, 1964
potato, 1964
potato psyllid, 1964
tomato apex necrosis virus,
Bemisia tabaci, 1559
tomato marchitez virus, 1559
tomato torrado virus, 1559
tomato marchitez virus,
Bemisia tabaci, 1559
tomato apex necrosis virus, 1559
tomato torrado virus, 1559
tomato spotted wilt virus,
Bemisia tabaci, 924
fitness, 924
nonvector herbivore, 924
tomato torrado virus,
Bemisia tabaci, 1559
tomato apex necrosis virus, 1559
tomato marchitez virus, 1559
Tospovirus,
Frankliniella fusca, 587
peanut, 587
resistant genotypes, 587
total hemocyte count,
biological control, 1582
median lethal dose, 1582
median lethal time, 1582
minimum inhibitory concentration, 1582
toxicant,
methyl eugenol, 727
monitoring traps, 727
polymer matrix, 727
spinosad, 727
toxic bait,
azadirachtin, 1772
Mediterranean fruit fly, 1772
neem, 1772
South American fruit fly, 1772
toxicity, 36, 131, 322, 970, 1788
Bacillus thuringiensis, 36
Bactrocera dorsalis, 970
bioactive compound, 131
Blattella asahinai, 322
carboxylesterases, 970
chitinase inhibitor, 1788
clove oil, 131
eastern subterranean termite, 1788
gamma radiation, 36
harborage preference, 322
kilogray, 36
mulch, 322
repellent, 131
SDS-PAGE, 36
Solenopsis invicta, 131
synergist, 970
toxicity risk,
biological control, 142
chemical control, 142
Trichogramma spp., 142
toxin,
Bacillus thuringiensis, 2467
dose–response assay, 2467
transgenic corn, 2467
western bean cutworm, 2467
trait hierarchy,
Rhopalosiphum padi, 1894
Schizaphis graminum, 1894
Sitobion avenae, 1894
specific resistance, 1894
transgenic,
Bacillus thuringiensis, 1025
Bt, 1025
Phthorimaea operculella, 1025
Solanum tuberosum, 1025
transgenic corn, 823, 1250, 2467
Bacillus thuringiensis, 1250, 2467
Diabrotica virgifera virgifera, 823
dose–response assay, 2467
Event MON 88017, 823
Event MON 89034, 823
Helicoverpa zea, 1250
pheromone trap, 1250
resistance management, 823
toxin, 2467
western bean cutworm, 2467
transgenic crop, 1659, 2473
Callosobruchus maculatus, 2473
cowpea, 2473
detection capacity, 1659
environmental risk assessment, 1659
insect, 1659
insect resistance management, 2473
simulation, 2473
statistical power, 1659
transgenic maize,
Cry1Ab/Cry2Aj protein, 2548
Cry1Ab/Vip3DA protein, 2548
insect resistance evaluation, 2548
Ostrinia furnacalis, 2548
translocation,
Acer platanoides, 874
distribution, 874
imidacloprid, 874
neonicotinoid, 874
transmission,
acquisition, 1878
life history, 1878
plant virus, 1878
resistance, 1878
trap, 1739, 2109
ethyl acetate, 1739
integrated pest management (IPM), 1739

kairomone, 1739
red palm weevil, 1739
Saffron Thread Yellow, 2109
Sunny Summer Yellow, 2109
Tartar Red, 2109
western cherry fruit fly, 2109
trap crop,
choice test, 283
cole crop, 283
harlequin bug, 283
Murgantia histrionica, 283
trapping,
Fusarium, 1218
integrated pest management (IPM), 1218
monitoring, 1218
vector, 1218
trap tree,
ambrosia beetle, 1780
bottle trap, 1780
forecasting activity, 1780
phenological indicator, 1780
treatment,
ant, 1545
control, 1545
eradication, 1545
invasive ants, 1545
tree resistance,
cortical parenchyma, 473
Phloeomyzus passerinii, 473
Populus x canadensis, 473
Tribolium castaneum, 513, 2247
Epimedium pubescens, 513
food-processing facility, 2247
heat treatment, 2247
Liposcelis bostrychophila, 513
model, 2247
mortality, 2247
(R)-carvone, 513
repellency, 513
Tribolium confusum,
essential oil, 1902
Juniperus oxycedrus ssp. *oxycedrus*, 1902
silica gel, 1902
Sitophilus oryzae, 1902
stored wheat, 1902
Trichogramma spp.,
biological control, 142
chemical control, 142
toxicity risk, 142
tritrophic interaction,
multivariate statistical analysis, 1053
principal component analysis, 1053
VOCs, 1053
Trogossitidae,
Cerambycidae, 206
Cleridae, 206
Histeridae, 206
Scolytinae, 206
tropical population,
cotton, 86
nonreproducing state, 86
Tropilaelaps mite,
honey bee, 1535
screening protocols, 1535
surveillance, 1535
Tsuga canadensis,
canopy translocation, 2399
hemlock woolly adelgid, 2399
neonicotinoid, 2399
systemic insecticide, 2399
T. urticae,
biological control, 80
Eotetranychus lewisi, 80
phytoseiids, 80
strawberry, 80
twospotted spider mite, 862, 1707, 2371
biological control, 2371
cotton, 862
fecundity, 2371
full-count, 1707
life table, 2371
mint, 1707
nontarget effect, 2371
presence-absence, 1707
Tetranychus urticae, 862
yield, 862

U

upper developmental threshold,
bootstrap percentile, 107
lower developmental threshold, 107
Performance-2 model, 107
sum of effective temperatures, 107
urban landscape,
Phenacoccus peruvianus, 1486
phenology, 1486
sampling, 1486
spatial distribution, 1486
urban pest,
invasive species, 311
species distribution model, 311
subterranean termite, 311

V

Vaccinium,
biological control, 73
Eulophidae, 73
parasitism, 73
Platygastridae, 73
value added traits,
defoliation, 830
Glycine max, 830
insect resistance, 830
Vapormate,
fumigation, 2341
lemon, 2341
orange, 2341
phytosanitary, 2341
variety,
mushroom, 115
nonlinear modeling, 115
thermal requirement, 115
Varroa destructor, 558, 566
amitraz, 558
grooming behavior, 566
honey bee, 558
mite fall, 566
miticides, 558
Russian honey bee, 566
tau-fluvalinate, 558
Varroa resistance, 566
Varroa resistance,
grooming behavior, 566
mite fall, 566
Russian honey bee, 566
Varroa destructor, 566
vector, 16, 1218
Fusarium, 1218
integrated pest management (IPM), 1218
leaf disk transmission assay, 16
monitoring, 1218
semipersistent, 16
trapping, 1218
vector control,
economic, 366
fitness costs, 366
insecticide resistance, 366
malaria, 366
verbenone,
antiaggregant, 221
lodgepole pine, 221
mountain pine beetle, 221
semiochemical, 221
verde plant bug,
cotton, 189
Creontiades signatus, 189
feeding injury, 189
Vienna-8,
chlorpyrifos, 1226
pesticide toxicity, 1226
spinosad, 1226
sterile insect technique, 1226
vinegar,
attractant compounds, 979
Musca domestica, 979
vineyard agroecosystem,
dispersal, 2296
Hemiptera vector, 2296
plant community, 2296
Vir-ex,
insect colony effect, 50
production, 50
SeMNPV, 50
Spexit, 50
vitamin,
insect infestation, 1058
mineral, 1058

Sitophilus granarius, 1058
storage, 1058
Vitis,
Acerophagus, 1716
Ferrisia gilli, 1716
grape, 1716
mealybug, 1716
VOCs,
multivariate statistical analysis, 1053
principal component analysis, 1053
tritrophic interaction, 1053
voltage-gate sodium channel,
F1538I mutation, 2457
fenpropathrin, 2457
molecular marker, 2457
Tetranychus cinnabarinus, 2457

W

walking,
DAMs, 2259
flight initiation, 2259
insecticide resistance, 2259
Washington state vineyard,
grape, 482
mealybug, 482
monitoring, 482
pheromone, 482
water volume,
integrated pest management (IPM), 2126
selective insecticide, 2126
systemics, 2126
4-way olfactometer,
habitat fidelity, 1932
one-choice, 1932
spotted wing drosophila, 1932
western bean cutworm, 1274, 2467
Bacillus thuringiensis, 2467
dose–response assay, 2467
economic injury level, 1274
economic threshold, 1274
larval survival, 1274
Striacosta albicosta, 1274
toxin, 2467
transgenic corn, 2467
western cherry fruit fly, 1511, 2109
apple maggot, 1511
Malus spectabilis, 1511
microsatellites, 1511
Rhagoletis indifferens, 1511
Saffron Thread Yellow, 2109
Sunny Summer Yellow, 2109
Tartar Red, 2109
trap, 2109
western conifer seed bug,
biological control, 633
egg-parasitoid, 633
host age, 633
western tarnished plant bug,
biophysical model, 124
degree-day, 124
temperature-dependent development, 124
wheat,
English grain aphid, 1473
germplasm, 1473
resistance, 1473
tolerance, 1473
wheat pests,
cross-resistance, 1029
native species, 1029
plant resistance, 1029
whitefly,
arthropod predators, 1260
biological control, 1260
cotton, 1260
plant stress, 1260
white grub, 159, 1112, 1456
botanical insecticides, 159
Cylas formicarius, 1456
Diabrotica, 1456
entomopathogenic nematode, 1112
essential oil, 159
GC-MS, 159
Heterorhabditis, 1112
Ipomoea, 1456
peanut, 1112
plant resistance, 1456
scarab larvae, 159
Steinernema, 1112
whitespotted sawyer,
CT scan, 1331
damage progression, 1331
fire-killed trees, 1331
salvage logging, 1331
wild bees,
agricultural landscape, 97
crop pollination, 97
seminatural areas pollinator habitat, 97
wine grape,
invasive carpophagous species, 2349
mortality, 2349
winery waste, 2349
winery waste,
invasive carpophagous species, 2349
mortality, 2349
wine grape, 2349
wing formation,
fecundity, 10
Nilaparvata lugens, 10
sublethal effect, 10
survival, 10
wireworm,
damage, 786
mortality, 786
potato, 786
recovery, 786
wood borers,
establishment, 1145
invasive, 1145
natural enemy introduction, 1145
parasitoid release, 1145
worker mortality,
acquisition, 277
Bombus terrestris, 277
Metarhizium anisopliae strain F52, 277
miniature dispenser, 277

X

Xyleborus glabratus,
avocado, 2093
phytosanitation, 2093
Raffaelea lauricola, 2093
redbay, 2093
Xylosandrus,
ornamental nursery, 289
plant-based essential oil, 289
Pyrethroid, 289

Y

yeast fermentation,
bed bug, 1802
carbon dioxide, 1802
chemical lure, 1802
pitfall, 1802
sugar, 1802
yield,
cotton, 862
Tetranychus urticae, 862
twospotted spider mite, 862

Z

Zea mays, 1240, 2241, 2354
behavior, 2354
borer damage, 2241
Diabrotica virgifera virgifera, 2354
fumonisin, 2241
Fusarium verticillioides, 2241
host recognition, 2354
insect-plant interaction, 1240
larva, 2354
N availability, 1240
Sesamia nonagrioides, 2241
Spodoptera exigua, 1240
Zebra chip,
Bactericera cockerelli, 1440
Candidatus Liberibacter solanacearum, 1440
LAMP, 1440
pathogen detection, 1440